



Igor Langer

Kontakt

Igor Langer

Publikationen (15)

Viehl C, Oertli D, Sauter G, Terracciano L, Langer I, Banz V, Ramser M, Rosenthal R, Dell-Kuster S, Gueller U, Weixler B, Zuber M. Presence of bone marrow micro-metastases in stage I-III colon cancer patients is associated with worse disease-free and overall survival. *Cancer Med* 2017; 6:918-927.

Langer I, Zuber M, Koechli O, Burger D, Infanger E, Wight E, Schnarwyler B, Bronz L, Viehl C, Hess T, Fehr M, Schaer G, Singer G, Berclaz G, Worni M, Gueller U, Swiss Multicenter Sentinel Lymph Node Study Group in Breast Cancer. Bone marrow micrometastases do not impact disease-free and overall survival in early stage sentinel lymph node-negative breast cancer patients. *Ann Surg Oncol* 2013; 21:401-7.

Viehl C, Gueller U, Langer I, Laffer U, Oertli D, Zuber M. Factors influencing the success of in vivo sentinel lymph node procedure in colon cancer patients: Swiss prospective, multicenter study sentinel lymph node procedure in colon cancer. *World J Surg* 2013; 37:873-7.

Gueller U, Zetti A, Worni M, Langer I, Cabalzar-Wondberg D, Viehl C, Demartines N, Zuber M. Molecular investigation of lymph nodes in colon cancer patients using one-step nucleic acid amplification (OSNA): a new road to better staging?. *Cancer* 2012; 118:6039-45.

Viehl C, Oertli D, Laffer U, Riehle H, Terracciano L, Ochsner A, Langer I, Cecini R, Gueller U, Zuber M. Sentinel lymph node procedure leads to upstaging of patients with resectable colon cancer: results of the Swiss prospective, multicenter study sentinel lymph node procedure in colon cancer. *Ann Surg Oncol* 2012; 19:1959-65.

Viehl C, Langer I, Gueller U, Zanetti-Dällenbach R, Moch H, Wight E, Oertli D, Zuber M. Prognostic impact and therapeutic implications of sentinel lymph node micro-metastases in early-stage breast cancer patients. *J Surg Oncol* 2011; 103:531-3.

Langer I, Gueller U, Viehl C, Moch H, Wight E, Harder F, Oertli D, Zuber M. Axillary lymph node dissection for sentinel lymph node micrometastases may be safely omitted in early-stage breast cancer patients: long-term outcomes of a prospective study. *Indian J Surg Oncol* 2010; 1:59-67.

Viehl C, Ochsner A, von Holzen U, Cecini R, Langer I, Gueller U, Laffer U, Oertli D, Zuber M. Inadequate quality of surveillance after curative surgery for colon cancer. *Ann Surg Oncol* 2010; 17:2663-9.

Langer I, Gueller U, Viehl C, Moch H, Wight E, Harder F, Oertli D, Zuber M. Axillary lymph node dissection for sentinel lymph node micrometastases may be safely omitted in early-stage breast cancer patients: long-term outcomes of a prospective study. *Ann Surg Oncol* 2009; 16:3366-74.

Zuber M, Viehl C, Gueller U, Langer I. [Follow-up recommendations for patients after curative treatment of invasive breast cancer]. *Ther Umsch* 2008; 65:319-21.

Langer I, Zuber M, Burger D, Infanger E, Uehlinger U, Wight E, Schnarwyler B, Bronz L, Oertli D, Hess T, Fehr M, Schaer G, Moch H, Koechli O, Berclaz G, Gueller U, Swiss Multicenter Study Group Sentinel Lymph Node in Breast Cancer. Accuracy of frozen section of sentinel lymph nodes: a prospective analysis of 659 breast cancer patients of the Swiss multicenter study. *Breast Cancer Res Treat* 2008; 113:129-36.

Langer I, Zuber M, Burger D, Infanger E, Uehlinger U, Wight E, Schnarwyler B, Bronz L, Oertli D, Hess T, Fehr M, Schaer G, Singer G, Berclaz G, Koechli O, Gueller U, Swiss Multicenter Sentinel Lymph Node Study Group in Breast Cancer. Association of the presence of bone marrow micrometastases with the sentinel lymph node status in 410 early stage breast cancer patients: results of the Swiss Multicenter Study. *Ann Surg Oncol* 2007; 14:1896-903.

Langer I, Burger D, Infanger E, Uehlinger U, Wight E, Schnarwyler B, Bronz L, Oertli D, Hess T, Fehr M, Schaer G, Koechli O, Berclaz G, Gueller U, Zuber M. Morbidity of sentinel lymph node biopsy (SLN) alone versus SLN and completion axillary lymph node dissection after breast cancer surgery: a prospective Swiss multicenter study on 659 patients. *Ann Surg* 2007; 245:452-61.

Langer I, Marti W, Gueller U, Moch H, Harder F, Oertli D, Zuber M. Axillary recurrence rate in breast cancer patients with negative sentinel lymph node (SLN) or SLN micrometastases: prospective analysis of 150 patients after SLN biopsy. *Ann Surg* 2005; 241:152-8.

Gueller U, Harder F, Oertli D, Marti W, Langer I, Moch H, Torhorst J, Viehl C, Schirp U, Nitzsche E, Zuber M. Selective axillary surgery in breast cancer patients based on positron emission tomography with 18F-fluoro-2-deoxy-D-glucose: not yet!. *Breast Cancer Res Treat* 2002; 71:171-3.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch