



Stefan Rahm

Kontakt

Stefan Rahm

Bereiche

Orthopädische Chirurgie und Traumatologie des Bewegungsapparates

Publikationen (11)

Rahm S, Hoch A, Tondelli T, Fuchs J, Zingg P. Revision rate of THA in patients younger than 40 years depends on primary diagnosis - a retrospective analysis with a minimum follow-up of 10 years. *Eur J Orthop Surg Traumatol* 2021; 31:1335-1344.

Rahm S, Jud L, Jungwirth-Weinberger A, Tondelli T, Falkowski A, Sutter R, Zingg P. Mid-term results after pinning and hip arthroscopy for mild slipped capital femoral epiphysis: a minimum five-year follow-up. *J Child Orthop* 2020; 14:521-528.

Kiarostami P, Dennler C, Roner S, Sutter R, Fürnstahl P, Farshad M, Rahm S, Zingg P. Augmented reality-guided periacetabular osteotomy-proof of concept. *J Orthop Surg Res* 2020; 15:540.

Jud L, Vlachopoulos L, Häller T, Fucentese S, Rahm S, Zingg P. The impact of mal-angulated femoral rotational osteotomies on mechanical leg axis: a computer simulation model. *BMC Musculoskelet Disord* 2020; 21:50.

Rahm S, Ebnetter L, Schultz B, Spross C, Külling F, Erhardt J. Comparison of interfragmentary compression in conventional and locked plating of proximal unicondylar tibia fractures : A biomechanical cadaver study. *Acta Orthop Belg* 2016; 82:586-592.

Spross C, Jost B, Rahm S, Winklhofer S, Erhardt J, Benninger E. How many radiographs are needed to detect angular stable head screw cut outs of the proximal humerus - A cadaver study. *Injury* 2014; 45:1557-63.

Gerber C, Rahm S, Catanzaro S, Farshad M, Moor B. Latissimus Dorsi Tendon Transfer for Treatment of Irreparable Posterosuperior Rotator Cuff Tears: Long-Term Results at a Minimum Follow-up of Ten Years. *J Bone Joint Surg Am* 2013; 95:1920-6.

Rahm S, Wieser K, Spross C, Vich M, Gerber C, Meyer D. Standard axillary radiographs of the shoulder may mimic posterior subluxation of the lateral end of the clavicle. *J Orthop Trauma* 2013; 27:622-6.

Meyer D, Rahm S, Farshad M, Lajtai G, Wieser K. Deltoid muscle shape analysis with magnetic resonance imaging in patients with chronic rotator cuff tears. *BMC Musculoskelet Disord* 2013; 14:247.

Rahm S, Spross C, Gerber F, Farshad M, Buck F, Espinosa N. Operative treatment of chronic irreparable achilles tendon ruptures with large flexor hallucis longus tendon transfers. *Foot Ankle Int* 2013; 34:1100-10.

Wieser K, Rahm S, Farshad M, Ek E, Gerber C, Meyer D. Stitch positioning influences the suture hold in supraspinatus tendon repair. *Knee Surg Sports Traumatol Arthrosc* 2012; 21:1587-92.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch