



Stephanie Rondeau

Kontakt

Stephanie Rondeau

Publikationen (8)

Moccia A, Rauch D, Vorobiof D, Lohri A, Ruegsegger C, Biaggi Rudolf C, Rusterholz C, Hayoz S, Ghielmini M, Raats J, Del Giglio A, Taverna C, Schär S, Vanazzi A, Rondeau S, Hitz F, Mingrone W, Pabst T, Cevreska L, Zucca E. Prolonged rituximab maintenance in follicular lymphoma patients: long-term results of the SAKK 35/03 randomized trial. *Blood Adv* 2020; 4:5951-5957.

Zucca E, Zander T, Bersvendsen H, Bargetzi M, Mingrone W, Krasniqi F, Dirnhofer S, Hayoz S, Hawle H, Vilei S, Ghielmini M, Kimby E, Novak U, Lohri A, Ferreri A, Rondeau S, Vanazzi A, Østenstad B, Mey U, Rauch D, Wahlin B, Hitz F, Hernberg M, Johansson A, de Nully Brown P, Hagberg H, Swiss Group for Clinical Cancer Research and the Nordic Lymphoma Group. Short regimen of rituximab plus lenalidomide in follicular lymphoma patients in need of first-line therapy. *Blood* 2019; 134:353-362.

Driessen C, Pabst T, Hitz F, Hawle H, Rondeau S, Berset C, Besse A, Besse L, Ribi K, Samaras P, Mey U, Rüfer A, Mach N, Betticher D, Cantoni N, Novak U, Müller R, Zander T. Promising activity of nelfinavir-bortezomib-dexamethasone (NeVd) in proteasome inhibitor-refractory multiple myeloma. *Blood* 2018

Ribi K, Rondeau S, Hitz F, Mey U, Enoiu M, Pabst T, Stathis A, Fischer N, Clough-Gorr K. Cancer-specific geriatric assessment and quality of life: important factors in caring for older patients with aggressive B-cell lymphoma. *Support Care Cancer* 2017; 25:2833-2842.

Hitz F, Rondeau S, Berardi Vilei S, Eckhardt K, Enoiu M, Rothermundt C, Schmidt A, Krasniqi F, Mach N, Caspar C, Samaras P, Cairoli A, Fischer N, Pabst T, Zucca E, Mey U. Rituximab, bendamustine and lenalidomide in patients with aggressive B-cell lymphoma not eligible for anthracycline-based therapy or intensive salvage chemotherapy - SAKK 38/08. *Br J Haematol* 2016

Taverna C, Zucca E, Heijnen I, Rusterholz C, Rondeau S, Biaggi Rudolf C, Lohri A, Vorobiof D, Rauch D, Raats J, Laszlo D, Vanazzi A, Del Giglio A, Cevreska L, Pabst T, Mingrone W, Hitz F, Martinelli G, Ghielmini M. Rituximab Maintenance for a Maximum of 5 Years After Single-Agent Rituximab Induction in Follicular Lymphoma: Results of the Randomized Controlled Phase III Trial SAKK 35/03. *J Clin Oncol* 2015; 34:495-500.

Zaman K, Rondeau S, Hawle H, Rudolf C, Perey L, Wilidiers H, Berset C, Mueller A, Rochlitz C, Hasler-Strub U, Mamot C, Winterhalder R, Neven P. Fulvestrant with or without selumetinib, a MEK 1/2 inhibitor, in breast cancer progressing after aromatase inhibitor therapy: a multicentre randomised placebo-controlled double-blind phase II trial, SAKK 21/08. *Eur J Cancer* 2015; 51:1212-20.

Lee C, Ashton-Key M, Cogliatti S, Rondeau S, Schmitz S, Ghielmini M, Cragg M, Johnson P. Expression of the inhibitory Fc gamma receptor IIB (FCGR2B, CD32B) on follicular lymphoma cells lowers the response rate to rituximab monotherapy (SAKK 35/98). *Br J Haematol* 2014; 168:145-8.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch