



Markus Borner

Kontakt

Markus Borner

Publikationen (7)

Pestalozzi B, Novak U, Nussbaum C, Seifert B, Bigler M, Bize V, Vilei S, Rageth C, Aebi S, Borner M, Buser K, Tausch C, Dedes K, Rochlitz C, Zimmermann S, von Moos R, Winterhalder R, Ruhstaller T, Mueller A. Adjuvant treatment recommendations for patients with ER-positive/HER2-negative early breast cancer by Swiss tumor boards using the 21-gene recurrence score (SAKK 26/10). *BMC cancer* 2017; 17:265.

Rochlitz C, Rordorf T, Rauch D, Müller A, Ruhstaller T, Vetter M, Trojan A, Hasler-Strub U, Cathomas R, Winterhalder R, Borner M, Bärtschi D, Bigler M, von Moos R, Bernhard J, Matter-Walstra K, Wicki A, Zaman K, Anchisi S, Küng M, Na K. SAKK 24/09: safety and tolerability of bevacizumab plus paclitaxel vs. bevacizumab plus metronomic cyclophosphamide and capecitabine as first-line therapy in patients with HER2-negative advanced stage breast cancer - a multicenter, randomized phase III trial. *BMC cancer* 2016; 16:780.

Templeton A, Ruhstaller T, Thürlimann B, Zaman K, von Moos R, Rochlitz C, Winterhalder R, von Rohr L, Müller A, Winkler A, Borner M, Dietrich D, Beyeler M, Hsu Schmitz S, Sun H, Surber C, Ribí K, Swiss Group for Clinical Cancer Research (SAKK) Coordinating Center. Prevention of palmar-plantar erythrodysesthesia with an antiperspirant in breast cancer patients treated with pegylated liposomal doxorubicin (SAKK 92/08). *Breast* 2014; 23:244-9.

Templeton A, Schiess R, Wild P, Rüschoff J, Thalmann G, Dietrich P, Aebersold R, Klingbiel D, Gillessen Sommer S, Müller B, Winterhalder R, Stenner F, Dutoit V, Cathomas R, Rothermundt C, Bärtschi D, Dröge C, Gautschi O, Borner M, Fechter E, Swiss Group for Clinical Cancer Research (SAKK). Phase 2 trial of single-agent everolimus in chemotherapy-naive patients with castration-resistant prostate cancer (SAKK 08/08). *Eur Urol* 2013; 64:150-8.

Ruhstaller T, Köberle D, von Briel C, Lombriser N, Lohri A, Pederiva S, von Moos R, Thierstein S, Borner M, Schuller J, Ribí K, Templeton A, Popescu R. Intense therapy in patients with locally advanced esophageal cancer beyond hope for surgical cure: a prospective, multicenter phase II trial of the Swiss Group for Clinical Cancer Research (SAKK 76/02). *Onkologie* 2010; 33:222-8.

Ribí K, Pestalozzi B, Lombriser N, Borner M, von Moos R, Moosmann P, Hess V, Roth A, Honegger H, Schuller J, Köberle D, Ruhstaller T. Is a change in patient-reported dysphagia after induction chemotherapy in locally advanced esophageal cancer a predictive factor for pathological response to neoadjuvant chemoradiation?. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer* 2009; 17:1109-1116.

Koeberle D, Herrmann R, Mora O, Ruhstaller T, Strasser F, Beretta K, Mingrone W, Caspar C, Dietrich D, Gerber D, Borner M, Saletti P, Swiss Group for Clinical Cancer Research. Patient-reported outcomes of patients with advanced biliary tract cancers receiving gemcitabine plus capecitabine: a multicenter, phase II trial of the Swiss Group for Clinical Cancer Research. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2008; 26:3702-8.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch