



Martin Soekler

Kontakt

Martin Soekler

Publikationen (5)

Kreissl S, Keller U, Kühnhardt D, Ostermann H, Hertenstein B, Aulitzky W, Maschmeyer G, Vieler T, Eich H, Baues C, Stein H, Fuchs M, Diehl V, Dietlein M, Engert A, Borchmann P, Bentz M, Hitz F, Topp M, Goergen H, Buehnen I, Kobe C, Moccia A, Greil R, Eichenauer D, Zijlstra J, Markova J, Meissner J, Feuring-Buske M, Soekler M, Beck H, Willenbacher W, Ludwig W, Pabst T, German Hodgkin Study Group. PET-guided eBEACOPP treatment of advanced-stage Hodgkin lymphoma (HD18): follow-up analysis of an international, open-label, randomised, phase 3 trial. *Lancet Haematol* 2021; 8:e398–e409.

Kampa-Schittenhelm K, Vogel W, Bonzheim I, Fend F, Horger M, Kanz L, Soekler M, Schittenhelm M. Dasatinib overrides the differentiation blockage in a patient with mutant- D816V positive CBF β -MYH11 leukemia. *Oncotarget* 2018; 9:11876–11882.

Borchmann P, Bentz M, Keller U, Kühnhardt D, Ostermann H, Schmitz N, Hertenstein B, Aulitzky W, Maschmeyer G, Vieler T, Eich H, Baues C, Stein H, Fuchs M, Kuhnert G, Diehl V, Dietlein M, Hitz F, Topp M, Goergen H, Kobe C, Lohri A, Greil R, Eichenauer D, Zijlstra J, Markova J, Meissner J, Feuring-Buske M, Hüttmann A, Dierlamm J, Soekler M, Beck H, Willenbacher W, Ludwig W, Pabst T, Engert A. PET-guided treatment in patients with advanced-stage Hodgkin's lymphoma (HD18): final results of an open-label, international, randomised phase 3 trial by the German Hodgkin Study Group. *Lancet* 2017; 390:2790–2802.

Kreissl S, Borchmann P, Engert A, Diehl V, Fuchs M, Flechtner H, Rueffer J, Shonukan O, Soekler M, Hitz F, Halbsguth T, Behringer K, Brillant C, Mayer A, Goergen H, Mueller H. Cancer-related fatigue in patients with and survivors of Hodgkin's lymphoma: a longitudinal study of the German Hodgkin Study Group. *Lancet Oncol* 2016; 17:1453–1462.

Behringer K, Lange E, Mahlberg R, Hassler A, Vogelhuber M, Hahn D, Mezger J, Krause S, Skoetz N, Böll B, von Tresckow B, Diehl V, Hallek M, Borchmann P, Stein H, Eich H, Engert A, Bentz M, Vieler T, Kriz J, Goergen H, Tanner A, Zijlstra J, Greil R, Markova J, Sasse S, Fuchs M, Topp M, Soekler M, Mathas S, Meissner J, Wilhelm M, Koch P, Lindemann H, Schalk E, Semrau R. Omission of dacarbazine or bleomycin, or both, from the ABVD regimen in treatment of early-stage favourable Hodgkin's lymphoma (GHSG HD13): an open-label, randomised, non-inferiority trial. *Lancet* 2014; 385:1418–27.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch