



R Winterhalder

Kontakt

R Winterhalder

Publikationen (7)

Glatzer M, Pless M, Pederiva S, Pestalozzi B, Berger M, Inauen R, Winterhalder R, Montemurro M, Horber D, Putora P. Choice of first line systemic treatment in pancreatic cancer among national experts. *Pancreatology* 2020; 20:686-690.

Jörger M, van Schaik R, Becker M, Hayoz S, Pollak M, Cathomas R, Winterhalder R, Gillessen Sommer S, Rothermundt C. Multidrug and toxin extrusion 1 and human organic cation transporter 1 polymorphisms in patients with castration-resistant prostate cancer receiving metformin (SAKK 08/09). *Prostate Cancer Prostatic Dis* 2015

Koerberle D, Kollar A, Saletti P, Roth A, Früh M, Kueng M, Popescu R, Schacher S, Hess V, Moosmann P, Anchisi S, Betticher D, von Moos R, Dietrich D, Brauchli P, Baertschi D, Matter K, Winterhalder R, Borner M, Herrmann R. Bevacizumab continuation versus no continuation after first-line chemotherapy plus bevacizumab in patients with metastatic colorectal cancer: a randomized phase III non-inferiority trial (SAKK 41/06). *Ann Oncol* 2015; 26:709-14.

Strebel R, Winterhalder R, Morant R, Pless M, U H, Fehr M, Gillessen S, Schmid H, Sulser T, Cathomas R. Multidisciplinary care in patients with prostate cancer: room for improvement. *Support Care Cancer* 2013; 21:2327-33.

Helbling D, Bougel S, Leibundgut E, Benhattar J, Yan P, Baertschi D, Borner M, Widmer L, Saletti P, Rauch D, Madlung A, Winterhalder R, Burkhard R, Gloor B, Bosman F, Sun H, Gautschi O, Bodoky G, Köberle D. Neoadjuvant chemoradiotherapy with or without panitumumab in patients with wild-type KRAS, locally advanced rectal cancer (LARC): a randomized, multicenter, phase II trial SAKK 41/07. *Ann Oncol* 2012; 24:718-25.

Rochlitz C, Zaman K, Mayer M, Cliffe B, Herrmann R, Mannhart-Harms M, Muller A, Rauch D, Winterhalder R, von Moos R, Schonenberger A, Fehr M, Bühlmann M, Suter T, Huober J, Spirig C, Lerch S, Ruhstaller T, Swiss Group for Clinical Cancer Research (SAKK). Combination of bevacizumab and 2-weekly pegylated liposomal doxorubicin as first-line therapy for locally recurrent or metastatic breast cancer. A multicenter, single-arm phase II trial (SAKK 24/06). *Ann Oncol* 2010; 22:80-5.

Köberle D, Rust C, Bieri G, Neuweiler J, Terraciano L, Ruhstaller T, Heitzmann F, Hess V, Winterhalder R, von Moos R, Burkhard R, Toepfer M. Phase II study of capecitabine and oxaliplatin given prior to and concurrently with preoperative pelvic radiotherapy in patients with locally advanced rectal cancer. *British journal of cancer* 2008; 98:1204-9.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch