



D Rauch

Kontakt

D Rauch

Publikationen (7)

Metaxas Y, von Moos R, Rauch D, Appenzeller C, Waibel C, Schmid S, Froesch P, Biaggi-Rudolf C, Perrino M, Maconi A, Schneider M, Mark M, Ceresoli G, Zucali P, Pless M, Grossi F, Eboulet E, Früh M, Swiss Group for Clinical Cancer Research (SAKK). Lurbinectedin as second- or third-line palliative therapy in malignant pleural mesothelioma: an international, multi-centre, single-arm, phase II trial (SAKK 17/16). *Ann Oncol* 2020; 31:495–500.

Helbling D, Bougel S, Leibundgut E, Benhantar J, Yan P, Baertschi D, Borner M, Widmer L, Saletti P, Rauch D, Madlung A, Winterhalder R, Burkhardt R, Gloor B, Bosman F, Sun H, Gautschi O, Bodoky G, Köberle D. Neoadjuvant chemoradiotherapy with or without panitumumab in patients with wild-type KRAS, locally advanced rectal cancer (LARC): a randomized, multicenter, phase II trial SAKK 41/07. *Ann Oncol* 2012; 24:718–25.

Rochlitz C, Zaman K, Mayer M, Cliffe B, Herrmann R, Mannhart-Harms M, Müller A, Rauch D, Winterhalder R, von Moos R, Schonberger A, Fehr M, Bühlmann M, Suter T, Huober J, Spirig C, Lerch S, Ruhstaller T, Swiss Group for Clinical Cancer Research (SAKK). Combination of bevacizumab and 2-weekly pegylated liposomal doxorubicin as first-line therapy for locally recurrent or metastatic breast cancer. A multicenter, single-arm phase II trial (SAKK 24/06). *Ann Oncol* 2010; 22:80–5.

Borner M, Mingrone W, Lanz D, Dietrich D, Kappeler A, Roth A, Ruhstaller T, Caspar C, Pestalozzi B, Helbling D, Trojan A, Hess V, Rauch D, Saletti P, von Moos R, Köberle D, Swiss Group for Clinical Cancer Research (SAKK), Bern, Switzerland. Adding cetuximab to capecitabine plus oxaliplatin (XELOX) in first-line treatment of metastatic colorectal cancer: a randomized phase II trial of the Swiss Group for Clinical Cancer Research SAKK. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2008; 19:1288–92.

Hess D, Ribi K, Ballabeni P, Schuller J, Rauch D, Rochlitz C, Mattmann S, Schonberger A, Pagani O, Thürlmann B, Köberle D, Swiss Group for Clinical Cancer Research. Capecitabine and vinorelbine as first-line treatment in elderly patients (> or =65 years) with metastatic breast cancer. A phase II trial (SAKK 25/99). *Oncology* 2007; 73:228–37.

Borner M, Roth A, Lanz D, Brauchli P, Honegger H, Köberle D, Herrmann R, Rauch D, Saletti P, Wernli M, Popescu R, Dietrich D, Bernhard J, Swiss Group for Clinical Cancer Research (SAKK), Berne, Switzerland. A randomized phase II trial of capecitabine and two different schedules of irinotecan in first-line treatment of metastatic colorectal cancer: efficacy, quality-of-life and toxicity. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2005; 16:282–8.

Hess D, Goldhirsch A, Castiglione-Gertsch M, Rufener B, Ballabeni P, Rauch D, Aebi S, Pagani O, Thürlmann B, Swiss Group of Clinical Cancer Research (SAKK). Capecitabine and vinorelbine in elderly patients (> or =65 years) with metastatic breast cancer: a phase I trial (SAKK 25/99). *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2004; 15:1760–5.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch