



D Dietrich

Kontakt

D Dietrich

Publikationen (8)

Koeberle D, Kollar A, Saletti P, Roth A, Früh M, Kueng M, Popescu R, Schacher S, Hess V, Moosmann P, Anchisi S, Betticher D, von Moos R, Dietrich D, Brauchli P, Baertschi D, Matter K, Winterhalder R, Borner M, Herrmann R. Bevacizumab continuation versus no continuation after first-line chemotherapy plus bevacizumab in patients with metastatic colorectal cancer: a randomized phase III non-inferiority trial (SAKK 41/06). *Ann Oncol* 2015; 26:709-14.

Borner M, Mingrone W, Lanz D, Dietrich D, Kappeler A, Roth A, Ruhstaller T, Caspar C, Pestalozzi B, Helbling D, Trojan A, Hess V, Rauch D, Saletti P, von Moos R, Koeberle D, Swiss Group for Clinical Cancer Research (SAKK), Bern, Switzerland. Adding cetuximab to capecitabine plus oxaliplatin (XELOX) in first-line treatment of metastatic colorectal cancer: a randomized phase II trial of the Swiss Group for Clinical Cancer Research SAKK. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2008; 19:1288-92.

Perey L, Goldhirsch A, Köberle D, Clément P, Dietrich D, Fiche M, Wildiers H, Nolé F, Zaman K, Hawle H, Paridaens R, Thürlimann B. Clinical benefit of fulvestrant in postmenopausal women with advanced breast cancer and primary or acquired resistance to aromatase inhibitors: final results of phase II Swiss Group for Clinical Cancer Research Trial (SAKK 21/00). *Ann Oncol* 2007; 18:64-9.

Leyvraz S, Pestalozzi B, Dietrich D, Cerny T, Wernli M, Sessa C, Fey M, Christinat A, Honegger H, Guillou L, Herrmann R, Swiss Group for Clinical Cancer Research (SAKK). Treatment of advanced soft-tissue sarcomas using a combined strategy of high-dose ifosfamide, high-dose doxorubicin and salvage therapies. *British journal of cancer* 2006; 95:1342-7.

Leyvraz S, Honegger H, Dietrich D, Fey M, Sessa C, Cerny T, Lissoni A, Jundt G, Zweifel M, Swiss Group for Clinical Cancer Research. Long-term results of a multicenter SAKK trial on high-dose ifosfamide and doxorubicin in advanced or metastatic gynecologic sarcomas. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2006; 17:646-51.

Borner M, Roth A, Lanz D, Brauchli P, Honegger H, Köberle D, Herrmann R, Rauch D, Saletti P, Wernli M, Popescu R, Dietrich D, Bernhard J, Swiss Group for Clinical Cancer Research (SAKK), Berne, Switzerland. A randomized phase II trial of capecitabine and two different schedules of irinotecan in first-line treatment of metastatic colorectal cancer: efficacy, quality-of-life and toxicity. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2005; 16:282-8.

Morant R, Hering F, Hanselmann S, Gschwend A, Wernli M, Cerny T, Rochlitz C, Bauer J, Borner M, Bonomo M, Gillissen Sommer S, Dietrich D, Bernhard J, Schmid H. Capecitabine in hormone-resistant metastatic prostatic carcinoma - a phase II trial. *British journal of cancer* 2004; 90:1312-7.

Waeber M, Castiglione-Gertsch M, Dietrich D, Thürlimann B, Goldhirsch A, Brunner K, Borner M, Swiss Group for Clinical Cancer Research (SAKK). Adjuvant therapy after excision and radiation of isolated postmastectomy locoregional breast cancer recurrence: definitive results of a phase III randomized trial (SAKK 23/82) comparing tamoxifen with observation. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2003; 14:1215-21.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch