



## Shu-Fang Hsu Schmitz

### Kontakt

Shu-Fang Hsu Schmitz

## Publikationen (7)

Lee C, Ashton-Key M, Cogliatti S, Rondeau S, Schmitz S, Ghielmini M, Cragg M, Johnson P. Expression of the inhibitory Fc gamma receptor IIB (FCGR2B, CD32B) on follicular lymphoma cells lowers the response rate to rituximab monotherapy (SAKK 35/98). *Br J Haematol* 2014; 168:145-8.

Martinelli G, Zucca E, Dietrich P, Lohri A, Vorobiof D, Heizmann M, Stahel R, Stupp R, Okkinga E, Bassi S, Hess U, Cerny T, Utiger U, Schmitz S, Ghielmini M. Long-term follow-up of patients with follicular lymphoma receiving single-agent rituximab at two different schedules in trial SAKK 35/98. *J Clin Oncol* 2010; 28:4480-4.

Strasser F, Demmer-Steingruber R, Böhme C, Schmitz S, Thürlimann B, Cerny T, Gillessen Sommer S. Prevention of docetaxel- or paclitaxel-associated taste alterations in cancer patients with oral glutamine: a randomized, placebo-controlled, double-blind study. *The Oncologist* 2008; 13:337-46.

Ghielmini M, Cerny T, Bargetzi M, Ketterer N, Stahel R, Pichert G, Schefer H, Betticher D, Fey M, Waltzer U, Bertoni F, Cogliatti S, Schmitz S, Swiss Group for Clinical Cancer Research. Effect of single-agent rituximab given at the standard schedule or as prolonged treatment in patients with mantle cell lymphoma: a study of the Swiss Group for Clinical Cancer Research (SAKK). *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2005; 23:705-11.

Sabbioni M, Hürny C, Goldhirsch A, Herrmann R, Perey L, Bonnefoi H, Thürlimann B, Gertsch M, Schmitz S, Siegrist H, Bernhard J, Swiss Group for Clinical Cancer Research (SAKK). Does subjective burden of early breast cancer and its treatment affect immune measures during adjuvant therapy?. *Breast Cancer Res Treat* 2004; 87:75-86.

Ghielmini M, Vorobiof D, Bargetzi M, Lohri A, Helg C, Kovacsovics T, Stupp R, Zucca E, Hess U, Peccatori F, Martinelli G, Betticher D, Fey M, Waltzer U, Hummerjohann J, Pichert G, Cogliatti S, Schmitz S, Cerny T. Prolonged treatment with rituximab in patients with follicular lymphoma significantly increases event-free survival and response duration compared with the standard weekly x 4 schedule. *Blood* 2004; 103:4416-23.

Bertoni F, Cotter F, Cavalli F, Zucca E, Auer R, Jones C, Baldini L, Ponzoni M, Bertolini F, Pichert G, Fey M, Cerny T, Ghielmini M, Schmitz S, Cogliatti S, Conconi A, Swiss Group for Clinical Cancer Research. Immunoglobulin heavy chain genes somatic hypermutations and chromosome 11q22-23 deletion in classic mantle cell lymphoma: a study of the Swiss Group for Clinical Cancer Research. *British journal of haematology* 2004; 124:289-98.

## Projekte (0)

Keine Resultate gefunden.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)