



**Martin Wetzke**

**Kontakt**

Martin Wetzke

## Publikationen (19)

Stallhofer J, Beigel F, Schnitzler F, Tillack-Schreiber C, Glas J, Lohse P, Wetzke M, Konrad-Zerna A, Friedrich M, Brand S. Lipocalin-2 Is a Disease Activity Marker in Inflammatory Bowel Disease Regulated by IL-17A, IL-22, and TNF- $\alpha$  and Modulated by IL23R Genotype Status. *Inflamm Bowel Dis* 2015; 21:2327-40.

Tillack C, Koburger M, Wagner J, Glas J, Diegelmann J, Koglin S, Dombrowski Y, Schaubert J, Wollenberg A, Maier H, Wetzke M, Ehmann L, Friedrich M, Laubender R, Papay P, Vogelsang H, Stallhofer J, Beigel F, Bedynek A, Brand S. Anti-TNF antibody-induced psoriasiform skin lesions in patients with inflammatory bowel disease are characterised by interferon- $\gamma$ -expressing Th1 cells and IL-17A/IL-22-expressing Th17 cells and respond to anti-IL-12/IL-23 antibody treatment. *Gut* 2013; 63:567-77.

Glas J, Czamara D, Diegelmann J, Göke B, Friedrich M, Steib C, Beigel F, Wetzke M, Tsekere E, Olszak T, Fries C, Stallhofer J, Bues S, Seiderer J, Brand S. IRGM variants and susceptibility to inflammatory bowel disease in the German population. *PloS one* 2013; 8:e54338.

Glas J, Duerr R, Franke A, Kamboh M, Achkar J, Balschun T, Müller-Myhsok B, Wolf C, Olszak T, Wetzke M, Diegelmann J, Pasciuto G, Czamara D, Seiderer J, Brand S. PTGER4 expression-modulating polymorphisms in the 5p13.1 region predispose to Crohn's disease and affect NF- $\kappa$ B and XBP1 binding sites. *PloS one* 2012; 7:e52873.

Glas J, Czamara D, Diegelmann J, Ochsenkühn T, Göke B, Wetzke M, Steib C, Stallhofer J, Beigel F, Friedrich M, Tillack C, Fries C, Olszak T, Wagner J, Seiderer J, Brand S. Analysis of IL12B gene variants in inflammatory bowel disease. *PloS one* 2012; 7:e34349.

Glas J, Czamara D, Diegelmann J, Karbalai N, Ochsenkühn T, Göke B, Steib C, Friedrich M, Stallhofer J, Tillack C, Beigel F, Wetzke M, Olszak T, Seiderer J, Wagner J, Brand S. PTPN2 gene variants are associated with susceptibility to both Crohn's disease and ulcerative colitis supporting a common genetic disease background. *PloS one* 2012; 7:e33682.

Glas J, Czamara D, Diegelmann J, Friedrich M, Steib C, Beigel F, Olszak T, Tillack C, Fries C, Wetzke M, Bayrle C, Seiderer J, Brand S. The role of osteopontin (OPN/SPP1) haplotypes in the susceptibility to Crohn's disease. *PloS one* 2011; 6:e29309.

Glas J, Czamara D, Diegelmann J, Müller-Myhsok B, Folwaczny M, Ochsenkühn T, Göke B, Weidinger M, Olszak T, Beigel F, Wetzke M, Pfennig S, Tengler B, Fischer D, Seiderer J, Brand S. Pregnane X receptor (PXR/NR112) gene haplotypes modulate susceptibility to inflammatory bowel disease. *Inflamm Bowel Dis* 2010; 17:1917-24.

Glas J, Seiderer J, Tillack C, Paschos E, Wetzke M, Diegelmann J, Czamara D, Brand S. Functional SFTPD gene variants are not associated with susceptibility to inflammatory bowel disease in the German population. *Inflamm Bowel Dis* 2010; 17:1439-40.

Glas J, Roeske D, Diegelmann J, Müller-Myhsok B, Ochsenkühn T, Göke B, Paschos E, Wetzke M, Pfennig S, Markus C, Seiderer J, Brand S. Role of PPARG gene variants in inflammatory bowel disease. *Inflamm Bowel Dis* 2010; 17:1057-8.

Jürgens M, Herrmann K, Lohse P, Göke B, Kreis M, Schnitzler F, Weidinger M, Beigel F, Tillack C, Pfennig S, Wagner J, Wetzke M, Glas J, Seiderer J, Laubender R, Brand S, Ochsenkühn T. The presence of fistulas and NOD2 homozygosity strongly predict intestinal stenosis in Crohn's disease independent of the IL23R genotype. *J Gastroenterol* 2010; 45:721-31.

Jürgens M, Ochsenkühn T, Glas J, Göke B, Lohse P, Tillack C, Schnitzler F, Stallhofer J, Pfennig S, Beigel F, Wetzke M, Wagner J, Seiderer J, Weidinger M, Hartl F, Laubender R, Brand S. Disease activity, ANCA, and IL23R genotype status determine early response to infliximab in patients with ulcerative colitis. *Am J Gastroenterol* 2010; 105:1811-9.

Török H, Müller-Myhsok B, Folwaczny C, Göke B, Folwaczny M, Ochsenkühn T, Lohse P, Klein W, Wetzke M, Teshome M, Tonenchi L, Endres I, Glas J, Brand S. Epistasis between Toll-like receptor-9 polymorphisms and variants in NOD2 and IL23R modulates susceptibility to Crohn's disease. *Am J Gastroenterol* 2009; 104:1723-33.

Glas J, Müller-Myhsok B, Ochsenkühn T, Göke B, Lohse P, Folwaczny M, Koletzko S, Lacher M, Schiemann U, Griga T, Epplen J, Klein W, Pfennig S, Wetzke M, Ripke S, Stallhofer J, Brand S. Novel genetic risk markers for ulcerative colitis in the IL2/IL21 region are in epistasis with IL23R and suggest a common genetic background for ulcerative colitis and celiac disease. *Am J Gastroenterol* 2009; 104:1737-44.

Glas J, Klein W, Epplen J, Schiemann U, Mussack T, Lohse P, Göke B, Ochsenkühn T, Folwaczny M, Müller-Myhsok B, Griga T, Jürgens M, Stallhofer J, Seiderer J, Pasciuto G, Tillack C, Diegelmann J, Pfennig S, Konrad A, Schmechel S, Wetzke M, Török H, Brand S. rs224136 on chromosome 10q21.1 and variants in PHOX2B, NCF4, and FAM92B are not major genetic risk factors for susceptibility to Crohn's disease in the German population. *Am J Gastroenterol* 2009; 104:665-72.

Schmechel S, Konrad A, Diegelmann J, Glas J, Wetzke M, Paschos E, Lohse P, Göke B, Brand S. Linking genetic susceptibility to Crohn's disease with Th17 cell function: IL-22 serum levels are increased in Crohn's disease and correlate with disease activity and IL23R genotype status. *Inflamm Bowel Dis* 2008; 14:204-12.

Glas J, Klein W, Epplen J, Folwaczny C, Lohse P, Göke B, Ochsenkühn T, Mussack T, Folwaczny M, Müller-Myhsok B, Griga T, Haller D, Pfennig S, Konrad A, Schmechel S, Dambacher J, Seiderer J, Schroff F, Wetzke M, Roeske D, Török H, Tonenchi L, Brand S. The ATG16L1 gene variants rs2241879 and rs2241880 (T300A) are strongly associated with susceptibility to Crohn's disease in the German population. *Am J Gastroenterol* 2007; 103:682-91.

Glas J, Epplen J, Schiemann U, Folwaczny C, Lohse P, Göke B, Ochsenkühn T, Müller-Myhsok B, Folwaczny M, Mussack T, Klein W, Griga T, Maier K, Seiderer J, Wetzke M, Konrad A, Török H, Schmechel S, Tonenchi L, Grassl C, Dambacher J, Pfennig S, Brand S. rs1004819 is the main disease-associated IL23R variant in German Crohn's disease patients: combined analysis of IL23R, CARD15, and OCTN1/2 variants. *PloS one* 2007; 2:e819.

Glas J, Folwaczny M, Ochsenkühn T, Brand S, Schnitzler F, Seiderer J, Lohse P, Schiemann U, Griga T, Epplen J, Klein W, Wetzke M, Mussack T, Müller-Myhsok B, Tonenchi L, Török H, Folwaczny C. Role of the NFKB1 -94ins/delATTG promoter polymorphism in IBD and potential interactions with polymorphisms in the CARD15/NOD2, IKBL, and IL-1RN genes. *Inflamm Bowel Dis* 2006; 12:606-11.

## Projekte (0)

Keine Resultate gefunden.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)