



**R Herrmann**

**Kontakt**

R Herrmann

## Publikationen (5)

Koeberle D, Kollar A, Saletti P, Roth A, Früh M, Kueng M, Popescu R, Schacher S, Hess V, Moosmann P, Anchisi S, Betticher D, von Moos R, Dietrich D, Brauchli P, Baertschi D, Matter K, Winterhalder R, Borner M, Herrmann R. Bevacizumab continuation versus no continuation after first-line chemotherapy plus bevacizumab in patients with metastatic colorectal cancer: a randomized phase III non-inferiority trial (SAKK 41/06). *Ann Oncol* 2015; 26:709-14.

Rochlitz C, Zaman K, Mayer M, Cliffe B, Herrmann R, Mannhart-Harms M, Muller A, Rauch D, Winterhalder R, von Moos R, Schonenberger A, Fehr M, Bühlmann M, Suter T, Huober J, Spirig C, Lerch S, Ruhstaller T, Swiss Group for Clinical Cancer Research (SAKK). Combination of bevacizumab and 2-weekly pegylated liposomal doxorubicin as first-line therapy for locally recurrent or metastatic breast cancer. A multicenter, single-arm phase II trial (SAKK 24/06). *Ann Oncol* 2010; 22:80-5.

Leyvraz S, Pestalozzi B, Dietrich D, Cerny T, Wernli M, Sessa C, Fey M, Christinat A, Honegger H, Guillou L, Herrmann R, Swiss Group for Clinical Cancer Research (SAKK). Treatment of advanced soft-tissue sarcomas using a combined strategy of high-dose ifosfamide, high-dose doxorubicin and salvage therapies. *British journal of cancer* 2006; 95:1342-7.

Borner M, Roth A, Lanz D, Brauchli P, Honegger H, Köberle D, Herrmann R, Rauch D, Saletti P, Wernli M, Popescu R, Dietrich D, Bernhard J, Swiss Group for Clinical Cancer Research (SAKK), Berne, Switzerland. A randomized phase II trial of capecitabine and two different schedules of irinotecan in first-line treatment of metastatic colorectal cancer: efficacy, quality-of-life and toxicity. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2005; 16:282-8.

Sabbioni M, Goldhirsch A, Herrmann R, Perey L, Bonnefoi H, Thürlimann B, Castiglione M, Bernhard J, Bacchi M, Siegrist H, Hürny C. Association between immunity and prognostic factors in early stage breast cancer patients before adjuvant treatment. *Breast cancer research and treatment* 2000; 59:279-87.

## Projekte (0)

Keine Resultate gefunden.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)