



## Michael T Hirschmann

### Kontakt

Michael T Hirschmann

## Publikationen (14)

Hess S, Moser L, Robertson E, Behrend H, Amsler F, Iordache E, Leclercq V, Hirschmann M. Osteoarthritic and non-osteoarthritic patients show comparable coronal knee joint line orientations in a cross-sectional study based on 3D reconstructed CT images. *Knee Surg Sports Traumatol Arthrosc* 2021

Hochreiter B, Moser L, Hess S, Hirschmann M, Amsler F, Behrend H. Osteoarthritic knees have a highly variable patellofemoral alignment: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2020

Hochreiter B, Hess S, Moser L, Hirschmann M, Amsler F, Behrend H. Correction to: Healthy knees have a highly variable patellofemoral alignment: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2019

Hochreiter B, Hess S, Moser L, Hirschmann M, Amsler F, Behrend H. Healthy knees have a highly variable patellofemoral alignment: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2019; 28:398–406.

Hess S, Moser L, Amsler F, Behrend H, Hirschmann M. Highly variable coronal tibial and femoral alignment in osteoarthritic knees: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2019; 27:1368–1377.

Hirschmann M, Moser L, Amsler F, Behrend H, Leclercq V, Hess S. Phenotyping the knee in young non-osteoarthritic knees shows a wide distribution of femoral and tibial coronal alignment. *Knee Surg Sports Traumatol Arthrosc* 2019; 27:1385–1393.

Hirschmann M, Moser L, Amsler F, Behrend H, Leclercq V, Hess S. Functional knee phenotypes: a novel classification for phenotyping the coronal lower limb alignment based on the native alignment in young non-osteoarthritic patients. *Knee Surg Sports Traumatol Arthrosc* 2019; 27:1394–1402.

Hirschmann M, Hess S, Behrend H, Amsler F, Leclercq V, Moser L. Phenotyping of hip–knee–ankle angle in young non-osteoarthritic knees provides better understanding of native alignment variability. *Knee Surg Sports Traumatol Arthrosc* 2019; 27:1378–1384.

Moser L, Hess S, Amsler F, Behrend H, Hirschmann M. Native non-osteoarthritic knees have a highly variable coronal alignment: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2019; 27:1359–1367.

Hochreiter B, Hirschmann M, Amsler F, Behrend H. Highly variable tibial tubercle–trochlear groove distance (TT–TG) in osteoarthritic knees should be considered when performing TKA. *Knee Surg Sports Traumatol Arthrosc* 2018

Lampart M, Behrend H, Moser L, Hirschmann M. Due to great variability fixed HKS angle for alignment of the distal cut leads to a significant error in coronal TKA orientation. *Knee Surg Sports Traumatol Arthrosc* 2018

Hirschmann M, Behrend H. Functional knee phenotypes: a call for a more personalised and individualised approach to total knee arthroplasty?. *Knee Surg Sports Traumatol Arthrosc* 2018

Forrer F, Hirschmann M, Rasch H. SPECT/CT for the assessment of painful knee prosthesis. *Nucl Med Commun* 2014; 35:782.

Rasch H, Falkowski A, Forrer F, Henckel J, Hirschmann M. 4D-SPECT/CT in orthopaedics: a new method of combined quantitative volumetric 3D analysis of SPECT/CT tracer uptake and component position measurements in patients after total knee arthroplasty. *Skeletal Radiol* 2013; 42:1215–23.

## Projekte (0)

Keine Resultate gefunden.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)