



Jochen Utikal

Kontakt

Jochen Utikal

Publikationen (5)

Váraljai R, Zimmer L, Al-Matary Y, Kaptein P, Albrecht L, Shannan B, Brase J, Gusenleitner D, Amaral T, Wyss N, Utikal J, Flatz L, Rambow F, Reinhardt H, Dick J, Engel D, Horn S, Ugurel S, Sondermann W, Livingstone E, Sucker A, Paschen A, Zhao F, Placke J, Klose J, Fendler W, Thommen D, Helfrich I, Schadendorf D, Roesch A. Author Correction: Interleukin 17 signaling supports clinical benefit of dual CTLA-4 and PD-1 checkpoint inhibition in melanoma. *Nat Cancer* 2023; 4:1395.

Váraljai R, Zimmer L, Al-Matary Y, Kaptein P, Albrecht L, Shannan B, Brase J, Gusenleitner D, Amaral T, Wyss N, Utikal J, Flatz L, Rambow F, Reinhardt H, Dick J, Engel D, Horn S, Ugurel S, Sondermann W, Livingstone E, Sucker A, Paschen A, Zhao F, Placke J, Klose J, Fendler W, Thommen D, Helfrich I, Schadendorf D, Roesch A. Interleukin 17 signaling supports clinical benefit of dual CTLA-4 and PD-1 checkpoint inhibition in melanoma. *Nat Cancer* 2023; 4:1292-1308.

Garbe C, Keim U, Amaral T, Berking C, Eigentler T, Flatz L, Gesierich A, Leiter U, Stadler R, Sunderkötter C, Tüting T, Utikal J, Wollina U, Zimmer L, Zouboulis C, Ascierto P, Eggermont A, Grob J, Hauschild A, Sekulovic L, Long G, Luke J, Michielin O, Peris K, Schadendorf D, Kirkwood J, Lorigan P, Central Malignant Melanoma Registry (CMMR). Prognosis of Patients With Primary Melanoma Stage I and II According to American Joint Committee on Cancer Version 8 Validated in Two Independent Cohorts: Implications for Adjuvant Treatment. *J Clin Oncol* 2022; 40:3741-3749.

Wagner N, Utikal J, Umansky V, Garbe C, Schadendorf D, Holland-Letz T, Herpel E, Sucker A, Lichtenberger R, Funder A, Kehrel C, Kemper C, Schuermans V, Tarnanidis K, Reith M, Gries M, Weide B, Gebhardt C. Tumor microenvironment-derived S100A8/A9 is a novel prognostic biomarker for advanced melanoma patients and during immunotherapy with anti-PD-1 antibodies. *J Immunother Cancer* 2019; 7:343.

Wagner N, Utikal J, Enk A, Umansky V, Garbe C, Naehrer H, Holland-Letz T, Busch C, Ikenberg K, Eubel J, Herpel E, Pflugfelder A, Lichtenberger R, Kehrel C, Tarnanidis K, Reith M, Weide B, Gebhardt C. Diminished levels of the soluble form of RAGE are related to poor survival in malignant melanoma. *Int J Cancer* 2015; 137:2607-17.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch