



Hans-Peter Brunner-La Rocca

Kontakt

Hans-Peter Brunner-La Rocca

Publikationen (26)

Simonavičius J, Maeder M, Eurlings C, Aizpurua A, Čelutkienė J, Barysienė J, Toggweiler S, Kaufmann B, Brunner-La Rocca H. Intensification of pharmacological decongestion but not the actual daily loop diuretic dose predicts worse chronic heart failure outcome: insights from TIME-CHF. *Clin Res Cardiol* 2020

Lanfear D, Li J, Brunner-La Rocca H, Kraus W, Sabbah H, Maeder M, Zeld N, Sanders-van Wijk S, Adams K, O'Connor C, Donahue M, Gui H, She R, Luzum J, Williams L. Polygenic Score for Beta-Blocker Survival Benefit in European Ancestry Patients with Reduced Ejection Fraction Heart Failure. *Circ Heart Fail* 2020

Stöhr R, Sanders-van Wijk S, Pfisterer M, Jeker U, Schuh A, Leibundgut G, Maeder M, Heine G, Brandenburg V, Brunner-La Rocca H. Limited role for fibroblast growth factor 23 in assessing prognosis in heart failure patients: data from the TIME-CHF trial. *Eur J Heart Fail* 2020

Simonavičius J, van Empel V, Knackstedt C, Purnaitė R, Čelutkienė J, Pfisterer M, Kaufmann B, Pfister O, Maeder M, Rickenbacher P, Sanders van-Wijk S, Brunner-La Rocca H. Prognostic Significance of Longitudinal Clinical Congestion Pattern in Chronic Heart Failure: Insights From TIME-CHF Trial. *Am J Med* 2019; 132:e679-e692.

Steinmann E, Brunner-La Rocca H, Maeder M, Kaufmann B, Pfisterer M, Rickenbacher P. Is the clinical presentation of chronic heart failure different in elderly versus younger patients and those with preserved versus reduced ejection fraction?. *Eur J Intern Med* 2018

Davarzani N, Brunner-La Rocca H, Peeters R, Pfisterer M, Zaugg C, Rolny V, Block D, de Boer R, Suter T, Karel J, Smirnov E, Rickenbacher P, Maeder M, Sanders-van Wijk S, TIME-CHF Investigators. Novel concept to guide systolic heart failure medication by repeated biomarker testing—results from TIME-CHF in context of predictive, preventive, and personalized medicine. *EPMA J* 2018; 9:161-173.

Rickenbacher P, Kaufmann B, Maeder M, Bernheim A, Goetschalckx K, Pfister O, Pfisterer M, Brunner-La Rocca H. Heart failure with mid-range ejection fraction: a distinct clinical entity? Insights from the Trial of Intensified versus standard Medical therapy in Elderly patients with Congestive Heart Failure (TIME-CHF). *Eur J Heart Fail* 2017

Davarzani N, Peeters R, Rickenbacher P, Pfisterer M, Gutmann M, Leibundgut G, Maeder M, Karel J, Sanders-van Wijk S, Brunner-La Rocca H. N-Terminal Pro-B-Type Natriuretic Peptide-Guided Therapy in Chronic Heart Failure Reduces Repeated Hospitalizations—Results From TIME-CHF. *J Card Fail* 2017; 23:382-389.

Uszko-Lencer N, Brunner-La Rocca H, Zugck C, Pfisterer M, Osswald S, Muzzarelli S, Gutmann M, Maeder M, Spruit M, Frankenstein L. Predicting hospitalization and mortality in patients with heart failure: The BARDICHE-index. *Int J Cardiol* 2016; 227:901-907.

Maeder M, Brunner-La Rocca H, Pfisterer M, Julius B, Yoon S, Suter T, Kiencke S, Tobler D, Rickli H, Zurek M. Prognostic Value of the Change in Heart Rate From the Supine to the Upright Position in Patients With Chronic Heart Failure. *J Am Heart Assoc* 2016; 5

Zurek M, Brunner-La Rocca H, Pfisterer M, Jeker U, Handschin R, Abbühl H, Sanders-van Wijk S, Muzzarelli S, Rickli H, Maeder M, TIME-CHF Investigators. Differential Prognostic Impact of Resting Heart Rate in Older Compared to Younger Patients with Chronic Heart Failure - Insights from TIME-CHF. *J Card Fail* 2015

Brunner-La Rocca H, Maeder M, Rickenbacher P, Tobler D, Pfisterer M, Krause F, Rolny V, Eurlings L, Knackstedt C, TIME-CHF investigators. Impact of worsening renal function related to medication in heart failure. *Eur J Heart Fail* 2014; 17:159-68.

van Empel V, Maeder M, Kiencke S, Pfisterer M, Muzzarelli S, Min S, Goetschalckx K, Bernheim A, Kaufmann B, Brunner-La Rocca H. Interaction between pulmonary hypertension and diastolic dysfunction in an elderly heart failure population. *J Card Fail* 2013; 20:98-104.

Sanders-van Wijk S, Brunner-La Rocca H, Pfisterer M, Peter M, Rickenbacher P, Erne P, Estlinbaum W, Rickli H, Nietlispach F, Maeder M, TIME-CHF Investigators. Long-term results of intensified, N-terminal-pro-B-type natriuretic peptide-guided versus symptom-guided treatment in elderly patients with heart failure: five-year follow-up from TIME-CHF. *Circ Heart Fail* 2013; 7:131-9.

Maeder M, Brunner-La Rocca H, Pfisterer M, Peter M, Vuillomenet A, Erne P, Gutmann M, Abbühl H, Rickli H, Rickenbacher P, TIME-CHF Investigators. N-terminal pro brain natriuretic peptide-guided management in patients with heart failure and preserved ejection fraction: findings from the Trial of Intensified versus standard medical therapy in elderly patients with congestive heart failure (TIME-CHF). *Eur J Heart Fail* 2013; 15:1148-56.

Huijts M, Brunner-La Rocca H, Pfisterer M, Schindler R, Maeder M, Muzzarelli S, Burkard T, Duits A, van Oostenbrugge R, TIME-CHF Investigators. Cognitive impairment in heart failure: results from the Trial of Intensified versus standard Medical therapy in Elderly patients with Congestive Heart Failure (TIME-CHF) randomized trial. *Eur J Heart Fail* 2013; 15:699-707.

Sanders-van Wijk S, Brunner-La Rocca H, Pfisterer M, Peter M, Vuillomenet A, Rickenbacher P, Erne P, Estlinbaum W, Rickli H, van Asselt A, TIME-CHF Investigators. Cost-effectiveness of N-terminal pro-B-type natriuretic-guided therapy in elderly heart failure patients: results from TIME-CHF (Trial of Intensified versus Standard Medical Therapy in Elderly Patients with Congestive Heart Failure). *JACC Heart Fail* 2013; 1:64-71.

Spruit M, Brunner-La Rocca H, Rickli H, Pfisterer M, Kiencke S, Uszko-Lencer N, Jeker U, Ammann P, Knackstedt C, Maeder M, TIME-CHF Investigators. Prognostic value of self-reported versus objectively measured functional capacity in patients with heart failure: results from the TIME-CHF (Trial of Intensified Versus Standard Medical Therapy in Elderly Patients with Congestive Heart Failure). *J Am Coll Cardiol* 2012; 60:2125-6.

Maeder M, Brunner-La Rocca H, Estlinbaum W, Kiencke S, Dieterle T, Weilenmann D, Hack D, Fehr T, Ammann P, Muzzarelli S, Pfisterer M, Rickli H, TIME-CHF Investigators. Incidence, clinical predictors, and prognostic impact of worsening renal function in elderly patients with chronic heart failure on intensive medical therapy. *Am Heart J* 2012; 163:407-14, 414.e1.

Muzzarelli S, Brunner-La Rocca H, Pfisterer M, Burkard T, Julius B, Nietlispach F, Rickli H, Toggweiler S, Maeder M, TIME-CHF Investigators. Frequency and predictors of hyperkalemia in patients >60 years of age with heart failure undergoing intense medical therapy. *Am J Cardiol* 2011; 109:693-8.

Brunner-La Rocca H, Rickli H, Pfisterer M, Osswald S, Pfister O, Leventhal M, Kiowski W, Jeker U, Maeder M, Schindler R, Muzzarelli S, Rickenbacher P, TIME-CHF Investigators. End-of-life preferences of elderly patients with chronic heart failure. *Eur Heart J* 2011; 33:752-9.

Kaiser C, Pfisterer M, Ricard I, Brunner-La Rocca H, De Servi S, Bonetti P, Bertel O, Hornig B, Pedrazzini G, Rickli H, Alber H, Eberli F, Erne P, Galatius S, BASKET-PROVE Study Group. Drug-eluting versus bare-metal stents in large coronary arteries. *N Engl J Med* 2010; 363:2310-9.

Maeder M, Thompson B, Brunner-La Rocca H, Kaye D. Hemodynamic basis of exercise limitation in patients with heart failure and normal ejection fraction. *J Am Coll Cardiol* 2010; 56:855-63.

Muzzarelli S, Brunner-La Rocca H, Pfisterer M, Buser P, Jeker U, Gutmann M, Handschin R, Rickli H, Maeder M, Leibundgut G, TIME-CHF Investigators. Predictors of early readmission or death in elderly patients with heart failure. *Am Heart J* 2010; 160:308-14.

Pfisterer M, Brunner-La Rocca H, Schindler R, Hilti P, Schieber M, Osterhues H, Suter T, Yoon S, Beer H, Dubach P, Jeker U, Vuillomenet A, Rickenbacher P, Erne P, Gutmann M, Rickli H, Buser P, TIME-CHF Investigators. BNP-guided vs symptom-guided heart failure therapy: the Trial of Intensified vs Standard Medical Therapy in Elderly Patients With Congestive Heart Failure (TIME-CHF) randomized trial. *JAMA : the journal of the American Medical Association* 2009; 301:383-92.

Maggiorini M, Brunner-La Rocca H, Peth S, Fischler M, Böhm T, Bernheim A, Kiencke S, Bloch K, Dehnert C, Naeije R, Lehmann T, Bärtsch P, Mairbörl H. Both tadalafil and dexamethasone may reduce the incidence of high-altitude pulmonary edema: a randomized trial. *Ann Intern Med* 2006; 145:497-506.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch