



Prof. Dr.med. Bruno Weder

Kontakt

Prof. Dr.med. Bruno Weder
Switzerland

Bereiche

Neurologie

Publikationen (37)

Missimer J, Habegger S, Pastore-Wapp M, Wiest R, Weder B. Sensing form – finger gaiting as key to tactile object exploration – a data glove analysis of a prototypical daily task. *Journal of NeuroEngineering and Rehabilitation* 2020; 17

Abela E, Missimer J, Pastore-Wapp M, Wiest R, Weder B. Early prediction of long-term tactile object recognition performance after sensorimotor stroke. *Cortex* 2019; 115:264-279.

Galovic M, Kägi G, Brugger F, Weber J, Mueller M, Vos S, Zbinden M, Pastore-Wapp M, Leisi N, Weder B. Diverging lesion and connectivity patterns influence early and late swallowing recovery after hemispheric stroke. *Hum Brain Mapp* 2017

Galovic M, Leisi N, Müller M, Weber J, Tettenborn B, Brugger F, Abela E, Weder B, Kägi G. Neuroanatomical correlates of tube dependency and impaired oral intake after hemispheric stroke. *Eur J Neurol* 2016; 23:926-34.

Kägi G, Leisi N, Galovic M, Müller-Baumberger M, Weder B. Prolonged impairment of deglutition in supratentorial ischaemic stroke: the predictive value of Parramatta Hospitals' Assessment of Dysphagia. *Swiss Med Wkly* 2016; 146:w14355.

Wiest R, Abela E, Missimer J, Schroth G, Hess C, Sturzenegger M, Wang D, Weder B, Federspiel A. Interhemispheric cerebral blood flow balance during recovery of motor hand function after ischemic stroke--a longitudinal MRI study using arterial spin labeling perfusion. *PLoS one* 2014; 9:e106327.

Strbian D, Engelter S, Davis S, Cordonnier C, Eskandari A, Parkkila A, Forss N, Weder B, Murao K, Meretoja A, Numminen H, Saver J, Seiffge D, Michel P, Tatlisumak T. Symptomatic intracranial hemorrhage after stroke thrombolysis: comparison of prediction scores. *Stroke* 2014; 45:752-8.

Galovic M, Leisi N, Müller M, Abela E, Kägi G, Weder B (2013). Clinical pattern of swallowing impairment influences recovery of dysphagia after stroke.

Galovic M, Leisi N, Müller M, Weber J, Abela E, Kägi G, Weder B. Lesion location predicts transient and extended risk of aspiration after supratentorial ischemic stroke. *Stroke* 2013; 44:2760-7.

Galovic M, Kägi G, Weder B (2013). Predictors of severe dysphagia in ischemic stroke patients. Präsentiert bei: Swiss Stroke Trialists' Meeting, St. Gallen

Galovic M, Leisi N, Müller M, Weber J, Abela E, Kägi G, Weder B (2013). Ischemic infarctions of frontal operculum significantly interfere with recovery from dysphagia. Präsentiert bei: Swiss Stroke Society Meeting, St. Gallen

Reker P, Weder B, Vehoff J (2012). Acute thrombosis of a large Developmental Venous Anomaly.

Reker P, Weder B, Weber J, Tettenborn B, Vehoff J (2012). Acute thrombosis of a large Developmental Venous Anomaly.

Galovic M, Leisi N, Müller M, Weber J, Abela E, Felbecker A, Kägi G, Weder B (2012). MRI-basierte Prädiktoren von Dysphagie nach akutem supratentoriellem ischämischem Hirninfarkt.

Galovic M, Leupold D, Kägi G, Weder B (2012). Characteristics of stroke mimics in intravenous thrombolysis.
Galovic M, Leisi N, Müller M, Weber J, Abela E, Felbecker A, Kägi G, Weder B (2012). MRI-based predictors of dysphagia after stroke. Präsentiert bei: 16th Congress of the European Federation of Neurological Societies, Stockholm, Sweden
Müller-Westermann J, Hengartner H, Weber J, Weder B, Felbecker A (2011). Secondary Moyamoya syndrome with multiple ischemic strokes in a 10-year old child from Central Africa with sickle cell disease.
Galovic M, Mittas S, Leupold D, Berger C, Siebel P, Kägi G, Weder B (2011). Lysescreening in einem Schlaganfallnetzwerk: Internet basierte Datenbank zur Überprüfung von Qualitätsindikatoren.
Sarikaya H, Mattle H, Ballinari P, Sztajzel R, Sekoranja L, Mueller F, Tettenborn B, Weder B, Odier C, Michel P, Lyrer P, Engelter S, Arnold M, Baumgartner R. Intravenous thrombolysis in nonagenarians with ischemic stroke. Stroke 2011; 42:1967-70.
Kägi G, Missimer J, Abela E, Seitz R, Weder B. Neural networks engaged in tactile object manipulation: patterns of expression among healthy individuals. Behav Brain Funct 2010; 6:71.
Engelter S, Nedeltchev K, Reichhart M, Mattle H, Tettenborn B, Hungerbühler H, Sztajzel R, Baumgartner R, Michel P, Arnold M, Lüthy R, Sarikaya H, Rutgers M, Hatz F, Georgiadis D, Fluri F, Sekoranja L, Schwegler G, Müller F, Weder B, Lyrer P. Intravenous thrombolysis in stroke attributable to cervical artery dissection. Stroke 2009; 40:3772-6.
Hömke L, Amunts K, Böning L, Fretz C, Binkofski F, Zilles K, Weder B. Analysis of lesions in patients with unilateral tactile agnosia using cytoarchitectonic probabilistic maps. Human brain mapping 2009; 30:1444-56.
Bohlhalter S, Abela E, Weniger D, Weder B. Impaired verbal memory in Parkinson disease: relationship to prefrontal dysfunction and somatosensory discrimination. Behav Brain Funct 2009; 5:49.
Weder B, Tettenborn B (2008). Die Geschichte der Klinik für Neurologie am Kantonsspital St. Gallen. Ein "kleines" Fach auf dem Weg zum grossen Fachgebiet.
Hartmann S, Missimer J, Stoeckel C, Abela E, Shah J, Seitz R, Weder B. Functional connectivity in tactile object discrimination: a principal component analysis of an event related fMRI-Study. PloS one 2008; 3:e3831.
Tettenborn B, Felbecker A, Weder B (2006). DRUG INTERACTION BETWEEN PHENPROCUMON AND VALPROATE: CLINICALLY RELEVANT INCREASE OF INR IN PATIENTS WITH ISCHEMIC STROKE AND SYMPTOMATIC SEIZURES. Präsentiert bei: 7th European Congress on Epileptology: Helsinki, Finland, 2-6 July 2006: Platform Session, Helsinki, Finland
Felbecker A, Erdelyi B, Hertle T, Böning L, Weder B. Involuntary movements during. S C H W E I Z E R A R C H I V F Ü R N E U R O L O G I E U N D P S Y C H I A T R I E 2006; 156(07):371-391
Felbecker A, Weder B, Tettenborn B (2006). Clinical Relevance of Drug Interactions between Valproate and Oral Anticoagulants.
Felbecker A, Weder B, Tettenborn B (2006). Clinical Relevance of Drug Interactions between Valproate and Oral Anticoagulants. Präsentiert bei: American Academy of Neurology (AAN) congress, San Diego

Kägi G, Weder B (2005). Volume Flow Rate Evaluation in Patients with Internal Carotid Artery Stenosis. Präsentiert bei: Meeting of the Swiss Cerebrovascular Group 2005, -

Felbecker A, Erdelyi B, Hertle T, Bönig L, Weder B (2005). Involuntary movements during yawning of an otherwise plegic arm in a patient with acute ischaemic stroke. Präsentiert bei: 176. Tagung der Schweizerischen Neurologischen Gesellschaft, St. Gallen

Felbecker A, Tettenborn B, Weder B. Clinically Relevant Drug Interaction Between Valproate and the Oral Anticoagulant Phenprocoumon - Klinisch relevante Interaktion zwischen Valproat und Phenprocoumon. Aktuelle Neurologie 2005; 2005; 32:288-290.

Felbecker A, Weder B, Tettenborn B. Clinical relevance of drug interaction between valproate and phenprocoumon. Epilepsia 2005; 46 (suppl. 6):267.

Felbecker A, Weder B, Tettenborn B. Klinisch relevante Interaktion zwischen Valproat und Phenprocoumon. Akt Neurol 2005; 32:288-290.

Felbecker A, Weder B, Tettenborn B. Valporate and phenprocoumon: Is there an interaction of clinical relevance?. Cardiovasc 2005; 6:13.

Felbecker A, Weder B, Tettenborn B. Clinically relevant rise of International Normalized Ratio (INR) values in patients with ischemic stroke and symptomatic seizures: drug interaction between phenprocoumon and valproate. Schweiz Arch Neurol Psych 2005; 156:379-379.

Kägi G, Weder B (2004). Axial myopathy and parkinsonism. Präsentiert bei: 174th Meeting of the Swiss Neurological Society, Geneva 2004, -

Projekte (4)

Clinical and MRI based predictors of prolonged dysphagia in ischemic stroke

Klinische Forschung - 01.01.2013 - 31.07.2014

Abgeschlossen

Pilot study on the effect of escitalopram on cortical plasticity and glutamate transmission in healthy subjects: A neurophysiological and Magnet Resonance Spectroscopy study

Klinische Forschung - 01.07.2011 - 31.05.2015

Automatisch geschlossen

Randomisierte, doppelblinde, placebokontrollierte klinische Prüfung der Phase III mit Parallelgruppen zur Beurteilung der Wirksamkeit und Sicherheit von Desmoteplase bei Patienten mit akuten ischämischen Schlaganfall DIAS III

Klinische Forschung - 12.03.2010 - 31.12.2012

Abgebrochen

The significance of sensory function for the recovery of hand paresis after ischemic stroke

Klinische Forschung - 07.01.2010 - 07.01.2010

Automatisch geschlossen