



## **Monica Castiglione-Gertsch**

### **Kontakt**

Monica Castiglione-Gertsch

## Publikationen (18)

Pagani O, Goldhirsch A, Coates A, Fey M, Collins J, Thürlimann B, Lindtner J, Holmberg S, Castiglione-Gertsch M, Gelber R, Price K, International Breast Cancer Study Group (IBCSG). Patterns of recurrence of early breast cancer according to estrogen receptor status: a therapeutic target for a quarter of a century. *Breast Cancer Res Treat* 2009; 117:319-24.

Pagani O, Goldhirsch A, Coates A, Forbes J, Murray E, Fey M, Thürlimann B, Lindtner J, Collins J, Crivellari D, Holmberg S, Gelber R, Price K, Castiglione-Gertsch M, Simoncini E, Gelber S, International Breast Cancer Study Group. Is adjuvant chemotherapy of benefit for postmenopausal women who receive endocrine treatment for highly endocrine-responsive, node-positive breast cancer? International Breast Cancer Study Group Trials VII and 12-93. *Breast Cancer Res Treat* 2009; 116:491-500.

Thürlimann B, Coates A, Castiglione-Gertsch M, Forbes J, Collins J, Colleoni M, Crivellari D, Holmberg S, Gelber R, Price K, Goldhirsch A. Is chemotherapy necessary for premenopausal women with lower-risk node-positive, endocrine responsive breast cancer? 10-year update of International Breast Cancer Study Group Trial 11-93. *Breast Cancer Res Treat* 2009; 113:137-44.

Viale G, Olszewski W, Knox F, Thürlimann B, Price K, Castiglione-Gertsch M, Gelber R, Gusterson B, Goldhirsch A, Orosz Z, Neven P, Öhlschlegel C, Giobbie-Hurder A, Regan M, Coates A, Mastropasqua M, Dell'Orto P, Maiorano E, MacGrogan G, Braye S, Breast International Group Trial 1-98. Prognostic and predictive value of centrally reviewed Ki-67 labeling index in postmenopausal women with endocrine-responsive breast cancer: results from Breast International Group Trial 1-98 comparing adjuvant tamoxifen with letrozole. *J Clin Oncol* 2008; 26:5569-75.

Pestalozzi B, Goldhirsch A, Coates A, Castiglione-Gertsch M, Viale G, Murray E, Thürlimann B, Snyder R, Lindtner J, Holmberg S, Gelber R, Price K, Gusterson B, Mallon E, Zahrieh D, International Breast Cancer Study Group. Distinct clinical and prognostic features of infiltrating lobular carcinoma of the breast: combined results of 15 International Breast Cancer Study Group clinical trials. *JCO* 2008; 26:3006-14.

Crivellari D, Chirgwin J, Smith I, Rabaglio M, Gladiëff L, Del Mastro L, Láng I, Colleoni M, Gelber R, Castiglione-Gertsch M, Paridaens R, Forbes J, Mauriac L, Mouridsen H, Thürlimann B, Price K, Coates A, Sun Z, Goldhirsch A. Letrozole compared with tamoxifen for elderly patients with endocrine-responsive early breast cancer: the BIG 1-98 trial. *JCO* 2008; 26:1972-9.

Viale G, Gusterson B, Goldhirsch A, Price K, Castiglione-Gertsch M, Gelber R, Rusca T, Grigolato P, Braye S, Öhlschlegel C, Pillay K, Kovács A, Brown R, Perin T, Golouh R, Mastropasqua M, Maiorano E, Regan M, Coates A. Chemoendocrine compared with endocrine adjuvant therapies for node-negative breast cancer: predictive value of centrally reviewed expression of estrogen and progesterone receptors--International Breast Cancer Study Group. *J Clin Oncol* 2008; 26:1404-10.

Viale G, Coates A, Goldhirsch A, Gelber R, Castiglione-Gertsch M, Gusterson B, Öhlschlegel C, Pillay K, Kovács A, Brown R, Perin T, Golouh R, Price K, Colleoni M, Maiorano E, Maffini F, Mastropasqua M, Regan M, International Breast Cancer Study Group. Predictive value of tumor Ki-67 expression in two randomized trials of adjuvant chemoendocrine therapy for node-negative breast cancer. *J Natl Cancer Inst* 2008; 100:207-12.

Rasmussen B, Altermatt H, Gelber R, Castiglione-Gertsch M, Goldhirsch A, Gusterson B, Thürlimann B, Coates A, Viale G, Braye S, Lacroix-Triki M, Regan M, Lykkesfeldt A, Dell'Orto P, Del Curto B, Henriksen K, Mastropasqua M, Price K, Méry E, BIG 1-98 Collaborative and International Breast Cancer Study Groups. Adjuvant letrozole versus tamoxifen according to centrally-assessed ERBB2 status for postmenopausal women with endocrine-responsive early breast cancer: supplementary results from the BIG 1-98 randomised trial. *The Lancet* 2008; 9:23-8.

Mouridsen H, Price K, Smith I, Colleoni M, Gelber R, Paridaens R, Forbes J, Mauriac L, Thürlimann B, Sun Z, Castiglione-Gertsch M, Rabaglio M, Coates A, Keshaviah A, Goldhirsch A. Cardiovascular adverse events during adjuvant endocrine therapy for early breast cancer using letrozole or tamoxifen: safety analysis of BIG 1-98 trial. *JCO* 2007; 25:5715-22.

Viale G, Gusterson B, Goldhirsch A, Price K, Castiglione-Gertsch M, Gelber R, Thürlimann B, Öhlschlegel C, Braye S, Orosz Z, Neven P, Raffoul J, Rasmussen B, Dell'Orto P, Mastropasqua M, Maiorano E, Regan M, Coates A. Prognostic and predictive value of centrally reviewed expression of estrogen and progesterone receptors in a randomized trial comparing letrozole and tamoxifen adjuvant therapy for postmenopausal early breast cancer: BIG 1-98. *J Clin Oncol* 2007; 25:3846-52.

International Breast Cancer Study Group, Coates A, Castiglione-Gertsch M, Price K, Gelber R, Pagani O, Simoncini E, Gelber S, Colleoni M, Goldhirsch A. Effects of a treatment gap during adjuvant chemotherapy in node-positive breast cancer: results of International Breast Cancer Study Group (IBCSG) Trials 13-93 and 14-93. *Ann Oncol* 2007; 18:1177-84.

Karlsson P, Goldhirsch A, Gelber R, Forbes J, Crivellari D, Thürlimann B, Fey M, Holmberg S, Collins J, Lindtner J, Murray E, Gusterson B, Castiglione-Gertsch M, Coates A, Price K, Cole B, Wallgren A. The role of the number of uninvolved lymph nodes in predicting locoregional recurrence in breast cancer. *JCO* 2007; 25:2019-26.

Coates A, Del Mastro L, Smith I, Chirgwin J, Nogaret J, Pienkowski T, Wardley A, Jakobsen E, Price K, Láng I, Colleoni M, Keshaviah A, Thürlimann B, Mouridsen H, Mauriac L, Forbes J, Paridaens R, Castiglione-Gertsch M, Gelber R, Goldhirsch A. Five years of letrozole compared with tamoxifen as initial adjuvant therapy for postmenopausal women with endocrine-responsive early breast cancer: update of study BIG 1-98. *JCO* 2007; 25:486-92.

Bernhard J, Coates A, Goldhirsch A, Price K, Thürlimann B, Aebi S, Collins J, Murray E, Forbes J, Gelber R, Hürrny C, Castiglione-Gertsch M, Zahrieh D, International Breast Cancer Study Group Trial VIII. Adjuvant chemotherapy followed by goserelin compared with either modality alone: the impact on amenorrhea, hot flashes, and quality of life in premenopausal patients--the International Breast Cancer Study Group Trial VIII. *JCO* 2007; 25:263-70.

Breast International Group (BIG) 1-98 Collaborative Group, Price K, Wardly A, Wardley A, Smith I, Rabaglio M, Gelber R, Castiglione-Gertsch M, Paridaens R, Forbes J, Mauriac L, Mouridsen H, Coates A, Keshaviah A, Thürlimann B, Goldhirsch A. A comparison of letrozole and tamoxifen in postmenopausal women with early breast cancer. *NEJM* 2005; 353:2747-57.

Gruber G, Goldhirsch A, Coates A, Gelber R, Fey M, Simoncini E, Thürlimann B, Carbone A, Crivellari D, Collins J, Golouh R, Lindtner J, Holmberg S, Rudenstam C, Castiglione-Gertsch M, Price K, Nasi M, Bonetti M, International Breast Cancer Study Group. Prognostic value of extracapsular tumor spread for locoregional control in premenopausal patients with node-positive breast cancer treated with classical cyclophosphamide, methotrexate, and fluorouracil: long-term observations from International Breast Cancer Study Group Trial VI. *JCO* 2005; 23:7089-97.

Colleoni M, Thürlimann B, Crivellari D, Murray E, Mendiola C, Pagani O, Castiglione-Gertsch M, Coates A, Price K, Fey M, Collins J, Zahrieh D, Gelber R, Holmberg S, Mattsson J, Rudenstam C, Lindtner J, Erzen D, Snyder R, Goldhirsch A. Site of primary tumor has a prognostic role in operable breast cancer: the international breast cancer study group experience. *JCO* 2005; 23:1390-400.

## Projekte (0)

Keine Resultate gefunden.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)