



Holger Eidtmann

Kontakt

Holger Eidtmann

Publikationen (26)

Loibl S, Untch M, von Minckwitz G, Mehta K, Jackisch C, Tiemann K, Fasching P, Pfitzner B, Eidtmann H, Hartmann A, Lederer B, Furlanetto J, Klimowicz A, Huober J, Darb-Esfahani S, Denkert C. Integrated Analysis of PTEN and p4EBP1 Protein Expression as Predictors for pCR in HER2-Positive Breast Cancer. *Clin Cancer Res* 2016; 22:2675-83.

Marme F, von Minckwitz G, Untch M, Paepke S, Loibl S, Kümmel S, Jackisch C, Huober J, Hilfrich J, Hanusch C, Gerber B, Eidtmann H, Denkert C, Costa S, Blohmer J, Lederer B, Schneeweiss A. Utility of the CPS+EG staging system in hormone receptor-positive, human epidermal growth factor receptor 2-negative breast cancer treated with neoadjuvant chemotherapy. *Eur J Cancer* 2015; 53:65-74.

Hein A, Ekici A, Hanusch C, Huober J, Liedtke C, Mau C, Moisse M, Müller V, Nekljudova V, Peuteman G, Rack B, Rübner M, Van Brussel T, Wang L, Weinshilboum R, Loibl S, Beckmann M, Kunz G, Lambrechts D, von Minckwitz G, Häberle L, Eidtmann H, Tesch H, Untch M, Hilfrich J, Schem C, Rezai M, Gerber B, Dan Costa S, Blohmer J, Schwedler K, Kittel K, Fehm T, Fasching P. Genetic variants in VEGF pathway genes in neoadjuvant breast cancer patients receiving bevacizumab: Results from the randomized phase III GeparQuinto study. *Int J Cancer* 2015; 137:2981-8.

von Minckwitz G, Thomssen C, Huober J, Denkert C, Alfer J, Jackisch C, Nekljudova V, Burchardi N, Loibl S, Simon E, Stickeler E, Potenberg J, Conrad B, Reimer T, Decker T, Eidtmann H, Eiermann W, Hackmann J, Möbus V, Marme F, German Breast Group Investigators. A randomized phase 2 study comparing EC or CMF versus nab-paclitaxel plus capecitabine as adjuvant chemotherapy for nonfrail elderly patients with moderate to high-risk early breast cancer (ICE II-GBG 52). *Cancer* 2015; 121:3639-48.

Loibl S, Mehta K, Nekljudova V, Blohmer J, Costa S, Denkert C, Eidtmann H, Gerber B, Hanusch C, Hilfrich J, Huober J, Schneeweiss A, Kümmel S, Paepke S, Untch M, Lederer B, Jackisch C, von Minckwitz G. Outcome after neoadjuvant chemotherapy in young breast cancer patients: a pooled analysis of individual patient data from eight prospectively randomized controlled trials. *Breast Cancer Res Treat* 2015; 152:377-87.

Hanusch C, Burchardi N, Engels K, Blohmer J, Costa S, Denkert C, Eidtmann H, Gerber B, Hilfrich J, Huober J, Jackisch C, Kümmel S, Paepke S, Untch M, Loibl S, Schneeweiss A, von Minckwitz G. Dual Blockade with Afatinib and Trastuzumab as Neoadjuvant Treatment for Patients with Locally Advanced or Operable Breast Cancer Receiving Taxane-Anthracycline Containing Chemotherapy-DAFNE (GBG-70). *Clin Cancer Res* 2015; 21:2924-31.

Fontanella C, Schneeweiss A, Paepke S, Jackisch C, Mehta K, Nekljudova V, Untch M, Neven P, von Minckwitz G, Huober J, Hilfrich J, Hanusch C, Lederer B, Gade S, Vanoppen M, Blohmer J, Costa S, Denkert C, Eidtmann H, Gerber B, Loibl S. Impact of body mass index on neoadjuvant treatment outcome: a pooled analysis of eight prospective neoadjuvant breast cancer trials. *Breast Cancer Res Treat* 2015; 150:127-39.

Loibl S, Fasching P, Andre F, Lindner J, Sotiriou C, Dykgers A, Guo S, Gade S, Nekljudova V, Loi S, Untch M, Pfitzner B, Huober J, Heinrichs C, von Minckwitz G, Schneeweiss A, Paepke S, Lehmann A, Rezai M, Zahm D, Sinn P, Khandan F, Eidtmann H, Dohnal K, Denkert C. PIK3CA mutations are associated with lower rates of pathologic complete response to anti-human epidermal growth factor receptor 2 (her2) therapy in primary HER2-overexpressing breast cancer. *J Clin Oncol* 2014; 32:3212-20.

von Minckwitz G, Huober J, Costa S, Tesch H, Hanusch C, Hilfrich J, Khandan F, Fasching P, Sinn B, Engels K, Mehta K, Nekljudova V, Klare P, Eidtmann H, Schneeweiss A, Loibl S, Salat C, Denkert C, Rezai M, Blohmer J, Jackisch C, Paepke S, Gerber B, Zahm D, Kümmel S, Untch M. Neoadjuvant carboplatin in patients with triple-negative and HER2-positive early breast cancer (GeparSixto; GBG 66): a randomised phase 2 trial. *Lancet Oncol* 2014; 15:747-56.

Gerber B, Loibl S, Nekljudova V, Hauschild M, Huober J, Blohmer J, Kunz G, Jackisch C, Solbach C, Hanusch C, Kittel K, Schrader I, Eggemann H, Tesch H, Fasching P, Rezai M, Eidtmann H, von Minckwitz G, Untch M. Surgical outcome after neoadjuvant chemotherapy and bevacizumab: results from the GeparQuinto study (GBG 44). *Ann Surg Oncol* 2014; 21:2517-24.

Loibl S, Denkert C, Hilfrich J, Konecny G, Fett W, Stickeler E, Harbeck N, Mehta K, Nekljudova V, von Minckwitz G, Huober J, Kümmel S, Volz C, Mau C, Blohmer J, Costa S, Eidtmann H, Fasching P, Gerber B, Hanusch C, Jackisch C, Untch M. Response and prognosis after neoadjuvant chemotherapy in 1,051 patients with infiltrating lobular breast carcinoma. *Breast Cancer Res Treat* 2014; 144:153-62.

von Minckwitz G, Mehta K, Zahm D, Untch M, Schneeweiss A, Paepke S, Kümmel S, Kaufmann M, Jackisch C, Huober J, Hilfrich J, Hanusch C, Gerber B, Eiermann W, Eidtmann H, Denkert C, Costa S, Blohmer J, Loibl S. Response-guided neoadjuvant chemotherapy for breast cancer. *J Clin Oncol* 2013; 31:3623-30.

von Minckwitz G, Fehm T, Barinoff J, Rüdiger T, Erbstoesser E, Fasching P, Karn T, Müller V, Jackisch C, Huober J, Hilfrich J, Tesch H, Schmitt W, Loibl S, Müller B, Blohmer J, Sinn B, Eidtmann H, Eiermann W, Gerber B, Denkert C. Ki67 measured after neoadjuvant chemotherapy for primary breast cancer. *Clin Cancer Res* 2013; 19:4521-31.

Lück H, Schuster T, Wollschläger K, Untch M, Thomssen C, Eidtmann H, Stickeler E, Richter B, Hubalek M, Jackisch C, Stähler A, Beckmann M, Heilmann V, Huober J, Schrader I, Loibl S, du Bois A, von Minckwitz G. Capecitabine plus paclitaxel versus epirubicin plus paclitaxel as first-line treatment for metastatic breast cancer: efficacy and safety results of a randomized, phase III trial by the AGO Breast Cancer Study Group. *Breast Cancer Res Treat* 2013; 139:779-87.

Huober J, Kümmel S, Mau C, Jackisch C, Khandan F, Costa S, Krabisch P, Loibl S, Nekljudova V, Untch M, Höß C, Gerber B, Fasching P, Hanusch C, Rezai M, Eidtmann H, Kittel K, Hilfrich J, Schwedler K, Blohmer J, Tesch H, Minckwitz G. Neoadjuvant chemotherapy with paclitaxel and everolimus in breast cancer patients with non-responsive tumours to epirubicin/cyclophosphamide (EC) ± bevacizumab - results of the randomised GeparQuinto study (GBG 44). *Eur J Cancer* 2013; 49:2284-93.

von Minckwitz G, Mehta K, Nekljudova V, Denkert C, Konecny G, Kaufmann M, Jackisch C, Huober J, Hilfrich J, Eiermann W, Gerber B, Fasching P, Eidtmann H, Costa S, Blohmer J, Untch M, Loibl S. Definition and impact of pathologic complete response on prognosis after neoadjuvant chemotherapy in various intrinsic breast cancer subtypes. *J Clin Oncol* 2012; 30:1796-804.

von Minckwitz G, Blohmer J, Huober J, Hauschild M, Fehm T, Müller B, Denkert C, Loibl S, Nekljudova V, Untch M, German Breast Group, Kunz G, Jackisch C, Gerber B, Eidtmann H, Rezai M, Fasching P, Tesch H, Eggemann H, Schrader I, Kittel K, Hanusch C, Kreienberg R, Solbach C, Arbeitsgemeinschaft Gynäkologische Onkologie-Breast Study Groups. Neoadjuvant chemotherapy and bevacizumab for HER2-negative breast cancer. *N Engl J Med* 2012; 366:299-309.

Untch M, Gerber B, Rezai M, Jackisch C, Huober J, Kühn T, Nekljudova V, von Minckwitz G, German Breast Group (GBG), Hanusch C, Tesch H, Kreienberg R, Loibl S, Bischoff J, Eidtmann H, Kaufmann M, Blohmer J, Hilfrich J, Strumberg D, Fasching P, Arbeitsgemeinschaft Gynäkologische Onkologie-Breast (AGO-B) Study Group. Lapatinib versus trastuzumab in combination with neoadjuvant anthracycline-taxane-based chemotherapy (GeparQuinto, GBG 44): a randomised phase 3 trial. *Lancet Oncol* 2012; 13:135-44.

von Minckwitz G, Untch M, Diebold K, Tiemann K, Hoffmann G, Henschen S, Schrader I, Hanusch C, Clemens M, Just M, Dietrich K, Eidtmann H, Holms F, Solbach C, Tesch H, Huober J, Loibl S, Darb-Esfahani S, Denkert C. Responsiveness of adjacent ductal carcinoma in situ and changes in HER2 status after neoadjuvant chemotherapy/trastuzumab treatment in early breast cancer-results from the GeparQuattro study (GBG 40). *Breast Cancer Res Treat* 2011

von Minckwitz G, Zahm D, Khandan F, Hoffmann G, Gerber B, Eidtmann H, Fend F, Dietel M, Mehta K, du Bois A, Blohmer J, Müller B, Loibl S, Budczies J, Hanusch C, Darb-Esfahani S, Hilfrich J, Weiss E, Huober J, Denkert C. Cytoplasmic poly(adenosine diphosphate-ribose) polymerase expression is predictive and prognostic in patients with breast cancer treated with neoadjuvant chemotherapy. *J Clin Oncol* 2011; 29:2150-7.

Riethdorf S, von Minckwitz G, Untch M, Eidtmann H, Tesch H, Holms F, Hilfrich J, Schrader I, Fehm T, Huober J, Roller M, Komor M, Loibl S, Rau T, Zhang L, Müller V, Pantel K. Detection and HER2 expression of circulating tumor cells: prospective monitoring in breast cancer patients treated in the neoadjuvant GeparQuattro trial. *Clin Cancer Res* 2010; 16:2634-45.

von Minckwitz G, Kühn T, du Bois A, Blohmer J, Thomssen C, Dan Costa S, Jackisch C, Kaufmann M, Mehta K, Hanusch C, Gerber B, Rezai M, Loibl S, Fasching P, Huober J, Tesch H, Bauerfeind I, Hilfrich J, Eidtmann H, Untch M. Capecitabine in addition to anthracycline- and taxane-based neoadjuvant treatment in patients with primary breast cancer: phase III GeparQuattro study. *J Clin Oncol* 2010; 28:2015-23.

Untch M, Kühn T, du Bois A, Blohmer J, Thomssen C, Dan Costa S, Jackisch C, Kaufmann M, Mehta K, Hanusch C, Gerber B, Rezai M, Loibl S, Fasching P, Huober J, Tesch H, Bauerfeind I, Hilfrich J, Eidtmann H, von Minckwitz G. Neoadjuvant treatment with trastuzumab in HER2-positive breast cancer: results from the GeparQuattro study. *J Clin Oncol* 2010; 28:2024-31.

Costa S, Tesch H, Gerber B, Baumann K, Thomssen C, Breitbach G, Ibishi S, Jackisch C, Mehta K, Weiss E, Ataseven B, Loibl S, Kaufmann M, Zahm D, Hilfrich J, Huober J, Eidtmann H, du Bois A, Blohmer J, von Minckwitz G. Neoadjuvant chemotherapy shows similar response in patients with inflammatory or locally advanced breast cancer when compared with operable breast cancer: a secondary analysis of the GeparTrio trial data. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2010; 28:83-91.

von Minckwitz G, Kaufmann M, Mehta K, Loibl S, Jackisch C, Costa S, Huober J, Gerber B, Hilfrich J, Eidtmann H, Hanusch C, Vogel P, Kümmel S, German Breast Group. Neoadjuvant vinorelbine-capecitabine versus docetaxel-doxorubicin-cyclophosphamide in early nonresponsive breast cancer: phase III randomized GeparTrio trial. *Journal of the National Cancer Institute* 2008; 100:542-51.

von Minckwitz G, Kaufmann M, Mehta K, Loibl S, Jackisch C, Costa S, Huober J, Gerber B, Hilfrich J, Eidtmann H, Hanusch C, Vogel P, Kümmel S, German Breast Group. Intensified neoadjuvant chemotherapy in early-responding breast cancer: phase III randomized GeparTrio study. *Journal of the National Cancer Institute* 2008; 100:552-62.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch