



P Brauchli

Kontakt

P Brauchli

Publikationen (8)

Ruhstaller T, Mingrone W, Montemurro M, Girschikofsky M, Schmidt S, Bitzer M, Bedenne L, Brauchli P, Stahl M, Hess V, Mariette C, Hawle H, Thuss-Patience P, Hayoz S, Schacher S, Knorrnschild J, Schnider A, Plasswilm L, Budach W, Eisterer W, Swiss Group for Clinical Cancer Research (SAKK), the German Esophageal Cancer Study Group, the Austrian 'Arbeitsgemeinschaft Medikamentöse Tumortherapie' (AGMT), and the Fédération Francophone de Cancérologie Digestive (FFCD)/Fédération de Recherche en Ch. Neoadjuvant chemotherapy followed by chemoradiation and surgery with and without cetuximab in patients with resectable esophageal cancer: a randomized, open-label, phase III trial (SAKK 75/08). *Ann Oncol* 2018; 29:1386-1393.

Klingbiel D, Thürlimann B, Brauchli P, von Moos R. Absence of Evidence is not Evidence of Absence: The Case of Non-Inferiority. *Ann Oncol* 2017

Pagani O, Rabaglio M, Aebi S, Ribi K, Rochlitz C, Rothgiesser K, Thürlimann B, Moos R, Zaman K, Goldhirsch A, Pestalozzi B, Munzone E, Klingbiel D, Ruhstaller T, Nolé F, Eppenberger S, Oehlschlegel C, Bernhard J, Brauchli P, Hess D, Mamot C. Do all patients with advanced HER2 positive breast cancer need upfront-chemo when receiving trastuzumab? Randomized Phase III trial SAKK 22/99. *Ann Oncol* 2016

Strasser F, Koeberle D, Kaasa S, Mauri S, Haefner M, Brauchli P, Klingbiel D, Hayoz S, Betticher D, Aebi S, Ribi K, Cathomas R, von Moos R, Blum D, Swiss Group for Clinical Cancer Research (SAKK). The effect of real-time electronic monitoring of patient-reported symptoms and clinical syndromes in outpatient workflow of medical oncologists: E-MOSAIC, a multicenter cluster-randomized phase III study (SAKK 95/06). *Ann Oncol* 2015

Koeberle D, Kollar A, Saletti P, Roth A, Früh M, Kueng M, Popescu R, Schacher S, Hess V, Moosmann P, Anchisi S, Betticher D, von Moos R, Dietrich D, Brauchli P, Baertschi D, Matter K, Winterhalder R, Borner M, Herrmann R. Bevacizumab continuation versus no continuation after first-line chemotherapy plus bevacizumab in patients with metastatic colorectal cancer: a randomized phase III non-inferiority trial (SAKK 41/06). *Ann Oncol* 2015; 26:709-14.

Jörger M, Zappa F, Klingbiel D, Brauchli P, Rothschild S, Gautschi O, Pless M, Ochsenbein A, von Moos R, Betticher D, Stahel R, Droege C, Früh M, Baty F, Brutsche M. Circulating microRNA profiling in patients with advanced non-squamous NSCLC receiving bevacizumab/erlotinib followed by platinum-based chemotherapy at progression (SAKK 19/05). *Lung Cancer* 2014; 85:306-13.

Zaman K, Aebi S, Brauchli P, Genton C, Berthod G, Giobbie-Hurder A, Simcock M, Lüthi J, Pagani O, Schonenberger A, Huober J, Thürlimann B, on behalf of the Swiss Group for Clinical Cancer Research (SAKK). Bone mineral density in breast cancer patients treated with adjuvant letrozole, tamoxifen, or sequences of letrozole and tamoxifen in the BIG 1-98 study (SAKK 21/07). *Ann Oncol* 2011

Borner M, Roth A, Lanz D, Brauchli P, Honegger H, Köberle D, Herrmann R, Rauch D, Saletti P, Wernli M, Popescu R, Dietrich D, Bernhard J, Swiss Group for Clinical Cancer Research (SAKK), Berne, Switzerland. A randomized phase II trial of capecitabine and two different schedules of irinotecan in first-line treatment of metastatic colorectal cancer: efficacy, quality-of-life and toxicity. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2005; 16:282-8.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch