



K Skarvan

Kontakt

K Skarvan

Publikationen (14)

Freiermuth D, Skarvan K, Filipovic M, Seeberger M, Bolliger D. Volatile anaesthetics and positive pressure ventilation reduce left atrial performance: a transthoracic echocardiographic study in young healthy adults. *Br J Anaesth* 2014; 112:1032-41.

Bolliger D, Seeberger M, Kasper J, Skarvan K, Seeberger E, Lurati Buse G, Buser P, Filipovic M. Remifentanyl does not impair left ventricular systolic and diastolic function in young healthy patients. *Br J Anaesth* 2011; 106:573-9.

Bolliger D, Seeberger M, Kasper J, Bernheim A, Schumann R, Skarvan K, Buser P, Filipovic M. Different effects of sevoflurane, desflurane, and isoflurane on early and late left ventricular diastolic function in young healthy adults. *Br J Anaesth* 2010; 104:547-54.

Wang J, Seeberger M, Skarvan K, Michaux I, Bernet F, Arsenic R, Buser P, Filipovic M. Intra-operative myocardial ischaemia cannot be detected by analysis of transmitral inflow patterns in patients undergoing off-pump coronary surgery. *European journal of anaesthesiology* 2008; 25:1-7.

Filipovic M, Michaux I, Wang J, Hunziker P, Skarvan K, Seeberger M. Effects of sevoflurane and propofol on left ventricular diastolic function in patients with pre-existing diastolic dysfunction. *British journal of anaesthesia* 2007; 98:12-8.

Filipovic M, Wang J, Michaux I, Hunziker P, Skarvan K, Seeberger M. Effects of halothane, sevoflurane and propofol on left ventricular diastolic function in humans during spontaneous and mechanical ventilation. *British journal of anaesthesia* 2005; 94:186-92.

Skarvan K, Filipovic M, Wang J, Brett W, Seeberger M. Use of myocardial tissue Doppler imaging for intraoperative monitoring of left ventricular function. *British journal of anaesthesia* 2003; 91:473-80.

Filipovic M, Seeberger M, Rohlf s R, Dergeloo O, Studer W, Atar D, Buser P, Skarvan K. Doppler indices of diastolic transmitral flow velocity are invalid indicators of myocardial ischaemia during high-dose dobutamine infusion in anaesthetized patients. *European journal of anaesthesiology* 2002; 19:789-95.

Skarvan K, Lambert A, Filipovic M, Seeberger M. Reference values for left ventricular function in subjects under general anaesthesia and controlled ventilation assessed by two-dimensional transoesophageal echocardiography. *European journal of anaesthesiology* 2001; 18:713-22.

Zwetsch G, Filipovic M, Skarvan K, Todorov A, Seeberger M. Transient recurrent laryngeal nerve palsy after failed placement of a transesophageal echocardiographic probe in an anesthetized patient. *Anesthesia and analgesia* 2001; 92:1422-3.

Seeberger M, Filipovic M, Rohlf s R, Dergeloo O, Studer W, Atar D, Buser P, Skarvan K. The diagnostic value of Doppler echocardiographic indexes of diastolic filling for detecting demand ischemia in anesthetized patients. *International journal of cardiac imaging* 2000; 16:437-46.

Steiner L, Filipovic M, Skarvan K, Seeberger M. Anaesthetists as echocardiographers can influence the perioperative management of patients with suspected aortic stenosis. *European journal of anaesthesiology* 2000; 17:463.

Filipovic M, Seeberger M, Schneider M, Schmid M, Pargger H, Hunziker P, Skarvan K. Transthoracic echocardiography for perioperative haemodynamic monitoring. *British journal of anaesthesia* 2000; 84:800-3.

Filipovic M, Seeberger M, Steiner L, Skarvan K. Transthoracic echocardiography pre-, intra-, and postoperatively. Canadian journal of anaesthesia = Journal canadien d'anesthésie 2000; 47:192.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch