



**Martin Bolli**

**Kontakt**

Martin Bolli

## Publikationen (10)

Kuemmerli C, Schneider M, Joliat G, Moeckli B, Ukegini K, Bolli M, Worni M, Birrer D. Trends in pancreatic surgery in Switzerland: a survey and nationwide analysis over two decades. *Langenbecks Arch Surg* 2022; 407:3423–3435.

Zumstein V, Vinzens F, Zettl A, Heinemann K, Koeberle D, von Flüe M, Bolli M. Systematic immunohistochemical screening for Lynch syndrome in colorectal cancer: a single centre experience of 486 patients. *Swiss Med Wkly* 2016; 146:w14315.

Nebiker C, Oertli D, von Holzen U, Adamina M, Muraro M, Mengus C, Zajac P, Sconocchia G, Zuber M, Tornillo L, Terracciano L, Bolli M, Rosso R, Han J, Eppenberger-Castori S, Iezzi G, Hirt C, Amicarella F, Cremonesi E, Huber X, Padovan E, Angrisani B, Droseser R, Spagnoli G. GM-CSF Production by Tumor Cells Is Associated with Improved Survival in Colorectal Cancer. *Clin Cancer Res* 2014; 20:3094–106.

Adamina M, Spagnoli G, Marti W, Oertli D, Heberer M, Jacob A, Huegli R, Bolli M, Viehl C, Frey D, Weber W, Rosenthal R, Zajac P. Intranodal immunization with a vaccinia virus encoding multiple antigenic epitopes and costimulatory molecules in metastatic melanoma. *Mol Ther* 2010; 18:651–9.

Rosenthal R, Heberer M, Oertli D, von Holzen U, Frey D, Weber W, Bolli M, Iezzi G, Zajac P, Feder-Mengus C, Adamina M, Bracci L, Groeper C, Spagnoli G. Differential responsiveness to IL-2, IL-7, and IL-15 common receptor gamma chain cytokines by antigen-specific peripheral blood naive or memory cytotoxic CD8+ T cells from healthy donors and melanoma patients. *J Immunother* 2009; 32:252–61.

Bolli M, Spagnoli G, Hudolin T, Terracciano L, Carafa V, Sanguedolce F, Feder C, Gueller U, Zajac P, Schultz-Thater E, Tornillo L. NY-ESO-1/LAGE-1 coexpression with MAGE-A cancer/testis antigens: a tissue microarray study. *Int J Cancer* 2005; 115:960–6.

von Holzen U, Heberer M, Oertli D, Marti W, Schumacher R, Feder C, Reschner A, Groeper C, Zajac P, Weber W, Bolli M, Adamina M, Spagnoli G. Selective responsiveness to common gamma chain cytokines in peripheral blood-derived cytotoxic T lymphocytes induced by Melan-A/MART-1(27–35)targeted active specific immunotherapy. *Int J Cancer* 2005; 115:248–55.

Spagnoli G, Adamina M, Bolli M, Weber W, Zajac P, Marti W, Oertli D, Heberer M, Harder F. Active antigen-specific immunotherapy of melanoma: from basic science to clinical investigation. *World J Surg* 2005; 29:692–9.

Schumacher R, Adamina M, Zurbriggen R, Bolli M, Padovan E, Zajac P, Heberer M, Spagnoli G. Influenza virosomes enhance class I restricted CTL induction through CD4+ T cell activation. *Vaccine* 2004; 22:714–23.

Bolli M, Sauter G, Heberer M, Harder F, Gambazzi F, Mirlacher M, Haas P, Dalquen P, Guller U, Adamina M, Kocher T, Spagnoli G. Tissue microarray evaluation of Melanoma antigen E (MAGE) tumor-associated antigen expression: potential indications for specific immunotherapy and prognostic relevance in squamous cell lung carcinoma. *Ann Surg* 2002; 236:785–93; discussion 793.

## Projekte (0)

Keine Resultate gefunden.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)