



Daniel Oertli

Kontakt

Daniel Oertli
Switzerland

Publikationen (35)

Zulewski H, Dettmer M, Minder A, Clerici T, Triponez F, Oertli D, Weidner S, Steinert H, Haldemann A, Christ E, Bilz S, Giovanella L, Komminoth P. Multidisciplinary approach for risk-oriented treatment of low-risk papillary thyroid cancer in Switzerland. *Swiss Med Wkly* 2019; 149:w14700.

Jakob M, Gueller U, Ochsner A, Oertli D, Zuber M, Viehl C. Lymph node ratio is inferior to pN-stage in predicting outcome in colon cancer patients with high numbers of analyzed lymph nodes. *BMC Surg* 2018; 18:81.

Viehl C, Oertli D, Sauter G, Terracciano L, Langer I, Banz V, Ramser M, Rosenthal R, Dell-Kuster S, Gueller U, Weixler B, Zuber M. Presence of bone marrow micro-metastases in stage I-III colon cancer patients is associated with worse disease-free and overall survival. *Cancer Med* 2017; 6:918-927.

Weixler B, Warschkow R, Ramser M, Droseler R, von Holzen U, Oertli D, Kettelhack C. Urgent surgery after emergency presentation for colorectal cancer has no impact on overall and disease-free survival: a propensity score analysis. *BMC Cancer* 2016; 16:208.

Larussin H, von Holzen U, Viehl C, Rezaeian F, Riehle H, Oertli D, Gueller U, Zuber M. Blue dye injection does not induce dissemination of epithelial cells during SLN procedure in colon cancer patients. *Int J Colorectal Dis* 2014; 29:689-92.

Nebiker C, Oertli D, von Holzen U, Adamina M, Muraro M, Mengus C, Zajac P, Sconocchia G, Zuber M, Tornillo L, Terracciano L, Bolli M, Rosso R, Han J, Eppenberger-Castori S, Iezzi G, Hirt C, Amicarella F, Cremonesi E, Huber X, Padovan E, Angrisani B, Droseler R, Spagnoli G. GM-CSF Production by Tumor Cells Is Associated with Improved Survival in Colorectal Cancer. *Clin Cancer Res* 2014; 20:3094-106.

Viehl C, Gueller U, Langer I, Laffer U, Oertli D, Zuber M. Factors influencing the success of in vivo sentinel lymph node procedure in colon cancer patients: Swiss prospective, multicenter study sentinel lymph node procedure in colon cancer. *World J Surg* 2013; 37:873-7.

Viehl C, Oertli D, Laffer U, Riehle H, Terracciano L, Ochsner A, Langer I, Cecini R, Gueller U, Zuber M. Sentinel lymph node procedure leads to upstaging of patients with resectable colon cancer: results of the Swiss prospective, multicenter study sentinel lymph node procedure in colon cancer. *Ann Surg Oncol* 2012; 19:1959-65.

Clerici T, Oertli D, Udelsman R. Minimally Invasive Techniques in Thyroid Surgery. In: *Surgery of the Thyroid and Parathyroid Glands*. 2nd Heidelberg: Springer, 2011. ISBN 3642234585. S. 175-186.

Viehl C, Langer I, Gueller U, Zanetti-Dällenbach R, Moch H, Wight E, Oertli D, Zuber M. Prognostic impact and therapeutic implications of sentinel lymph node micro-metastases in early-stage breast cancer patients. *J Surg Oncol* 2011; 103:531-3.

Langer I, Gueller U, Viehl C, Moch H, Wight E, Harder F, Oertli D, Zuber M. Axillary lymph node dissection for sentinel lymph node micrometastases may be safely omitted in early-stage breast cancer patients: long-term outcomes of a prospective study. *Indian J Surg Oncol* 2010; 1:59-67.

Businger A, Gueller U, Oertli D. Effect of the 50-hour workweek limitation on training of surgical residents in Switzerland. *Arch Surg* 2010; 145:558-63.

Viehl C, Ochsner A, von Holzen U, Cecini R, Langer I, Gueller U, Laffer U, Oertli D, Zuber M. Inadequate quality of surveillance after curative surgery for colon cancer. *Ann Surg Oncol* 2010; 17:2663-9.

Adamina M, Spagnoli G, Marti W, Oertli D, Heberer M, Jacob A, Huegli R, Bolli M, Viehl C, Frey D, Weber W, Rosenthal R, Zajac P. Intranodal immunization with a vaccinia virus encoding multiple antigenic epitopes and costimulatory molecules in metastatic melanoma. *Mol Ther* 2010; 18:651-9.

Langer I, Gueller U, Viehl C, Moch H, Wight E, Harder F, Oertli D, Zuber M. Axillary lymph node dissection for sentinel lymph node micrometastases may be safely omitted in early-stage breast cancer patients: long-term outcomes of a prospective study. *Ann Surg Oncol* 2009; 16:3366-74.

Rosenthal R, Heberer M, Oertli D, von Holzen U, Frey D, Weber W, Bolli M, Iezzi G, Zajac P, Feder-Mengus C, Adamina M, Bracci L, Groepper C, Spagnoli G. Differential responsiveness to IL-2, IL-7, and IL-15 common receptor gamma chain cytokines by antigen-specific peripheral blood naive or memory cytotoxic CD8+ T cells from healthy donors and melanoma patients. *J Immunother* 2009; 32:252-61.

Langer I, Zuber M, Burger D, Infanger E, Uehlinger U, Wight E, Schnarwyler B, Bronz L, Oertli D, Hess T, Fehr M, Schaer G, Moch H, Koechli O, Berclaz G, Gueller U, Swiss Multicenter Study Group Sentinel Lymph Node in Breast Cancer. Accuracy of frozen section of sentinel lymph nodes: a prospective analysis of 659 breast cancer patients of the Swiss multicenter study. *Breast Cancer Res Treat* 2008; 113:129-36.

Adamina M, Heberer M, Zuber M, Oertli D, Frey D, Gueller U, Zajac P, Schumacher R, Rosenthal R, Weber W, Spagnoli G. Heterologous prime-boost immunotherapy of melanoma patients with Influenza virosomes, and recombinant Vaccinia virus encoding 5 melanoma epitopes and 3 co-stimulatory molecules. A multi-centre phase I/II open labeled clinical trial. *Contemp Clin Trials* 2007; 29:165-81.

Langer I, Zuber M, Burger D, Infanger E, Uehlinger U, Wight E, Schnarwyler B, Bronz L, Oertli D, Hess T, Fehr M, Schaer G, Singer G, Berclaz G, Koechli O, Gueller U, Swiss Multicenter Sentinel Lymph Node Study Group in Breast Cancer. Association of the presence of bone marrow micrometastases with the sentinel lymph node status in 410 early stage breast cancer patients: results of the Swiss Multicenter Study. *Ann Surg Oncol* 2007; 14:1896-903.

Weber W, Gueller U, Jain N, Pietrobon R, Oertli D. Impact of surgeon and hospital caseload on the likelihood of performing laparoscopic vs open sigmoid resection for diverticular disease: a study based on 55,949 patients. *Arch Surg* 2007; 142:253-9; discussion 259.

Langer I, Burger D, Infanger E, Uehlinger U, Wight E, Schnarwyler B, Bronz L, Oertli D, Hess T, Fehr M, Schaer G, Koechli O, Berclaz G, Gueller U, Zuber M. Morbidity of sentinel lymph node biopsy (SLN) alone versus SLN and completion axillary lymph node dissection after breast cancer surgery: a prospective Swiss multicenter study on 659 patients. *Ann Surg* 2007; 245:452-61.

Viehl C, Gueller U, Hamel C, Riehle H, Plaass C, Marti W, Oertli D, Zuber M. Carbon dye staining of sentinel lymph nodes facilitates microstaging of colon cancer patients. *World J Surg* 2006; 30:453-6.

Adamina M, Oertli D, Spagnoli G, Amacker M, Zurbriggen R, Feder C, Groepper C, Rosenthal R, Weber W, Zajac P, Schumacher R, Heberer M. Advanced liposomal vectors as cancer vaccines in melanoma immunotherapy. *J Liposome Res* 2006; 16:195-204.

Gueller U, Turek J, Eubanks S, Delong E, Oertli D, Feldman J. Detecting pheochromocytoma: defining the most sensitive test. *Ann Surg* 2006; 243:102-7.

Gueller U, Safford S, Pietrobon R, Heberer M, Oertli D, Jain N. High hospital volume is associated with better outcomes for breast cancer surgery: analysis of 233,247 patients. *World J Surg* 2005; 29:994-9; discussion 999-1000.

von Holzen U, Heberer M, Oertli D, Marti W, Schumacher R, Feder C, Reschner A, Groeper C, Zajac P, Weber W, Bolli M, Adamina M, Spagnoli G. Selective responsiveness to common gamma chain cytokines in peripheral blood-derived cytotoxic T lymphocytes induced by Melan-A/MART-1(27-35)targeted active specific immunotherapy. *Int J Cancer* 2005; 115:248-55.

Spagnoli G, Adamina M, Bolli M, Weber W, Zajac P, Marti W, Oertli D, Heberer M, Harder F. Active antigen-specific immunotherapy of melanoma: from basic science to clinical investigation. *World J Surg* 2005; 29:692-9.

Gueller U, Oertli D. Sample size matters: a guide for surgeons. *World J Surg* 2005; 29:601-5.

Adamina M, Oertli D. Antigen specific active immunotherapy: lessons from the first decade. *Swiss Med Wkly* 2005; 135:212-21.

Langer I, Marti W, Gueller U, Moch H, Harder F, Oertli D, Zuber M. Axillary recurrence rate in breast cancer patients with negative sentinel lymph node (SLN) or SLN micrometastases: prospective analysis of 150 patients after SLN biopsy. *Ann Surg* 2005; 241:152-8.

Gueller U, Jain N, Curtis L, Oertli D, Heberer M, Pietrobon R. Insurance status and race represent independent predictors of undergoing laparoscopic surgery for appendicitis: secondary data analysis of 145,546 patients. *J Am Coll Surg* 2004; 199:567-75; discussion 575-7.

Gueller U, Heberer M, Harder F, Metzger U, Maurer R, Oertli D, Spagnoli G, Zuber M, Vorburger S, Bösch B, Schnider A, Zajac P, Marti W. Disseminated single tumor cells as detected by real-time quantitative polymerase chain reaction represent a prognostic factor in patients undergoing surgery for colorectal cancer. *Ann Surg* 2002; 236:768-75; discussion 775-6.

Oertli D, Heberer M, Harder F, Schumacher R, Adamina M, Padovan E, Kocher T, Noppen C, Zajac P, Marti W, Spagnoli G. Rapid induction of specific cytotoxic T lymphocytes against melanoma-associated antigens by a recombinant vaccinia virus vector expressing multiple immunodominant epitopes and costimulatory molecules in vivo. *Hum Gene Ther* 2002; 13:569-75.

Spagnoli G, Zajac P, Marti W, Oertli D, Padovan E, Noppen C, Kocher T, Adamina M, Heberer M. Cytotoxic T-cell induction in metastatic melanoma patients undergoing recombinant vaccinia virus-based immuno-gene therapy. *Recent Results Cancer Res* 2002; 160:195-201.

Gueller U, Harder F, Oertli D, Marti W, Langer I, Moch H, Torhorst J, Viehl C, Schirp U, Nitzsche E, Zuber M. Selective axillary surgery in breast cancer patients based on positron emission tomography with 18F-fluoro-2-deoxy-D-glucose: not yet!. *Breast Cancer Res Treat* 2002; 71:171-3.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen Rorschacher Strasse 95 CH-9007 St.Gallen T: +41 71 494 11 11 support.forschung@kssg.ch