



Felix Harder

Kontakt

Felix Harder

Publikationen (9)

Langer I, Gueller U, Viehl C, Moch H, Wight E, Harder F, Oertli D, Zuber M. Axillary lymph node dissection for sentinel lymph node micrometastases may be safely omitted in early-stage breast cancer patients: long-term outcomes of a prospective study. *Indian J Surg Oncol* 2010; 1:59-67.

Langer I, Gueller U, Viehl C, Moch H, Wight E, Harder F, Oertli D, Zuber M. Axillary lymph node dissection for sentinel lymph node micrometastases may be safely omitted in early-stage breast cancer patients: long-term outcomes of a prospective study. *Ann Surg Oncol* 2009; 16:3366-74.

Spagnoli G, Adamina M, Bolli M, Weber W, Zajac P, Marti W, Oertli D, Heberer M, Harder F. Active antigen-specific immunotherapy of melanoma: from basic science to clinical investigation. *World J Surg* 2005; 29:692-9.

Langer I, Marti W, Gueller U, Moch H, Harder F, Oertli D, Zuber M. Axillary recurrence rate in breast cancer patients with negative sentinel lymph node (SLN) or SLN micrometastases: prospective analysis of 150 patients after SLN biopsy. *Ann Surg* 2005; 241:152-8.

Viehl C, Harder F, Spichtin H, Terracciano L, Stammberger U, Eisner L, Gueller U, Marti W, Hamel C, Zuber M. Identification of sentinel lymph nodes in colon cancer depends on the amount of dye injected relative to tumor size. *World J Surg* 2003; 27:1285-90.

Bolli M, Sauter G, Heberer M, Harder F, Gambazzi F, Mirlacher M, Haas P, Dalquen P, Guller U, Adamina M, Kocher T, Spagnoli G. Tissue microarray evaluation of Melanoma antigen E (MAGE) tumor-associated antigen expression: potential indications for specific immunotherapy and prognostic relevance in squamous cell lung carcinoma. *Ann Surg* 2002; 236:785-93; discussion 793.

Gueller U, Heberer M, Harder F, Metzger U, Maurer R, Oertli D, Spagnoli G, Zuber M, Vorburger S, Bösch B, Schnider A, Zajac P, Marti W. Disseminated single tumor cells as detected by real-time quantitative polymerase chain reaction represent a prognostic factor in patients undergoing surgery for colorectal cancer. *Ann Surg* 2002; 236:768-75; discussion 775-6.

Oertli D, Heberer M, Harder F, Schumacher R, Adamina M, Padovan E, Kocher T, Noppen C, Zajac P, Marti W, Spagnoli G. Rapid induction of specific cytotoxic T lymphocytes against melanoma-associated antigens by a recombinant vaccinia virus vector expressing multiple immunodominant epitopes and costimulatory molecules in vivo. *Hum Gene Ther* 2002; 13:569-75.

Gueller U, Harder F, Oertli D, Marti W, Langer I, Moch H, Torhorst J, Viehl C, Schirp U, Nitzsche E, Zuber M. Selective axillary surgery in breast cancer patients based on positron emission tomography with 18F-fluoro-2-deoxy-D-glucose: not yet!. *Breast Cancer Res Treat* 2002; 71:171-3.

Projekte (0)

Keine Resultate gefunden.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch