



Bent Ejlertsen

Contact

Bent Ejlertsen

Publications (9)

Leone J, Vaz-Luis I, Colleoni M, Di Leo A, Goldhirsch A, Wardley A, Bonnefoi H, Láng I, Neven P, Harvey V, Ejlertsen B, Gelber R, Giobbie-Hurder A, Rabaglio M, Coates A, Thürlimann B, Regan M, Cole B, Lin N. Clinical behavior of recurrent hormone receptor-positive breast cancer by adjuvant endocrine therapy within the Breast International Group 1-98 clinical trial. *Cancer* 2020

Rabaglio M, Thürlimann B, Colleoni M, Goldhirsch A, Gelber R, Coates A, Castiglione M, Ruepp B, Wardley A, Bonnefoi H, Láng I, Neven P, Harvey V, Ejlertsen B, Giobbie-Hurder A, Maibach R, Sun Z, Regan M. Cumulative incidence of cardiovascular events under tamoxifen and letrozole alone and in sequence: a report from the BIG 1-98 trial. *Breast Cancer Res Treat* 2020

Ruhstaller T, Simoncini E, Gianni L, Rochlitz C, Kralidis E, Zaman K, Jassem J, Piccart-Gebhart M, Di Leo A, Gelber R, Coates A, Goldhirsch A, Thürlimann B, Regan M, Veyret C, Del Mastro L, Giobbie-Hurder A, Colleoni M, Jensen M, Ejlertsen B, de Azambuja E, Neven P, Láng I, Jakobsen E, Gladieff L, Bonnefoi H, Harvey V, Spazzapan S, Tondini C, members of the BIG 1-98 Collaborative Group and the International Breast Cancer Study Group. Adjuvant Letrozole and Tamoxifen Alone or Sequentially for Postmenopausal Women With Hormone Receptor-Positive Breast Cancer: Long-Term Follow-Up of the BIG 1-98 Trial. *J Clin Oncol* 2018;JCO1800440.

Borgquist S, Regan M, Gelber R, Price K, Rabaglio M, Goldhirsch A, Coates A, Smith I, von Moos R, Ejlertsen B, Debled M, Láng I, Colleoni M, Garber J, Ahern T, Giobbie-Hurder A, Thürlimann B. Cholesterol, Cholesterol-Lowering Medication Use, and Breast Cancer Outcome in the BIG 1-98 Study. *J Clin Oncol* 2017; 35:1179-1188.

Chirgwin J, Colleoni M, Láng I, Neven P, Forbes J, Rabaglio M, Smith I, Goldhirsch A, Gelber R, Debled M, Ejlertsen B, Price K, Coates A, Giobbie-Hurder A, Thürlimann B. Treatment Adherence and Its Impact on Disease-Free Survival in the Breast International Group 1-98 Trial of Tamoxifen and Letrozole, Alone and in Sequence. *J Clin Oncol* 2016; 34:2452-9.

Metzger Filho O, Coates A, Regan M, Price K, Debled M, Ejlertsen B, Colleoni M, Gelber R, Thürlimann B, Winer E, Viale G, Gusterson B, Mallon E, Giobbie-Hurder A, Goldhirsch A. Relative Effectiveness of Letrozole Compared With Tamoxifen for Patients With Lobular Carcinoma in the BIG 1-98 Trial. *J Clin Oncol* 2015; 33:2772-9.

Ewertz M, Goldhirsch A, Coates A, Smith I, Láng I, Colleoni M, Gelber R, Rabaglio M, Paridaens R, Forbes J, Bonnefoi H, Thürlimann B, Price K, Ejlertsen B, Regan M, Gray K, Mouridsen H. Obesity and risk of recurrence or death after adjuvant endocrine therapy with letrozole or tamoxifen in the breast international group 1-98 trial. *J Clin Oncol* 2012; 30:3967-75.

Regan M, BIG 1-98 Collaborative Group, Thürlimann B, Coates A, Gelber R, Price K, Rabaglio M, Wardley A, Láng I, Smith I, Forbes J, Mauriac L, Ejlertsen B, Goldhirsch A, Giobbie-Hurder A, Neven P, International Breast Cancer Study Group (IBCSG). Assessment of letrozole and tamoxifen alone and in sequence for postmenopausal women with steroid hormone receptor-positive breast cancer: the BIG 1-98 randomised clinical trial at 8.1 years median follow-up. *Lancet Oncol* 2011; 12:1101-8.

Chirgwin J, Coates A, Goldhirsch A, Regan M, Bonnefoi H, Ejlertsen B, Thürlimann B, Price K, Smith I, Sun Z, BIG 1-98 Collaborative and International Breast Cancer Study Groups. The advantage of letrozole over tamoxifen in the BIG 1-98 trial is consistent in younger postmenopausal women and in those with chemotherapy-induced menopause. *Breast Cancer Res Treat* 2011; 131:295-306.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch