



Richard Greil

Contact

Richard Greil

Publications (14)

Bauer T, Santoro A, Lin C, Garrido-Laguna I, Jörger M, Greil R, Spreafico A, Yau T, Goebeler M, Hütter-Krönke M, Perotti A, Juif P, Lu D, Barys L, Cremasco V, Pelletier M, Evans H, Fabre C, Doi T. Phase I/Ib, open-label, multicenter, dose-escalation study of the anti-TGF- β monoclonal antibody, NIS793, in combination with spartalizumab in adult patients with advanced tumors. *J Immunother Cancer* 2023; 11

Harrington K, Burtness B, Greil R, Soulières D, Tahara M, de Castro G, Psyrrri A, Brana I, Basté N, Neupane P, Bratland Å, Fuereder T, Hughes B, Mesia R, Ngamphaiboon N, Rordorf T, Wan Ishak W, Lin J, Gumuscu B, Swaby R, Rischin D. Pembrolizumab With or Without Chemotherapy in Recurrent or Metastatic Head and Neck Squamous Cell Carcinoma: Updated Results of the Phase III KEYNOTE-048 Study. *J Clin Oncol* 2022;JCO2102508.

Rischin D, Harrington K, Greil R, Soulières D, Tahara M, de Castro G, Psyrrri A, Brana I, Neupane P, Bratland Å, Fuereder T, Hughes B, Mesia R, Ngamphaiboon N, Rordorf T, Ishak W, Hong R, Mendoza R, Jia L, Chirovsky D, Norquist J, Jin F, Burtness B. Pembrolizumab alone or with chemotherapy for recurrent or metastatic head and neck squamous cell carcinoma: Health-related quality-of-life results from KEYNOTE-048. *Oral Oncol* 2022; 128:105815.

Burtness B, Rischin D, Greil R, Soulières D, Tahara M, de Castro G, Psyrrri A, Brana I, Basté N, Neupane P, Bratland Å, Fuereder T, Hughes B, Mesia R, Ngamphaiboon N, Rordorf T, Wan Ishak W, Ge J, Swaby R, Gumuscu B, Harrington K. Pembrolizumab Alone or With Chemotherapy for Recurrent/Metastatic Head and Neck Squamous Cell Carcinoma in KEYNOTE-048: Subgroup Analysis by Programmed Death Ligand-1 Combined Positive Score. *J Clin Oncol* 2022; 40:2321-2332.

Kreissl S, Keller U, Kühnhardt D, Ostermann H, Hertenstein B, Aulitzky W, Maschmeyer G, Vieler T, Eich H, Baues C, Stein H, Fuchs M, Diehl V, Dietlein M, Engert A, Borchmann P, Bentz M, Hitz F, Topp M, Goergen H, Buehnen I, Kobe C, Moccia A, Greil R, Eichenauer D, Zijlstra J, Markova J, Meissner J, Feuring-Buske M, Soekler M, Beck H, Willenbacher W, Ludwig W, Pabst T, German Hodgkin Study Group. PET-guided eBEACOPP treatment of advanced-stage Hodgkin lymphoma (HD18): follow-up analysis of an international, open-label, randomised, phase 3 trial. *Lancet Haematol* 2021; 8:e398-e409.

Burtness B, Harrington K, Greil R, Soulières D, Tahara M, de Castro G, Psyrrri A, Basté N, Neupane P, Bratland Å, Fuereder T, Hughes B, Mesia R, Ngamphaiboon N, Rordorf T, Wan Ishak W, Hong R, González Mendoza R, Roy A, Zhang Y, Gumuscu B, Cheng J, Jin F, Rischin D. Pembrolizumab alone or with chemotherapy versus cetuximab with chemotherapy for recurrent or metastatic squamous cell carcinoma of the head and neck (KEYNOTE-048): a randomised, open-label, phase 3 study. *Lancet* 2019; 394:1915-1928.

Scheithauer W, Putora P, Grünberger B, Eisterer W, Wöll E, Prager G, Schaberl-Moser R, Greil R, Glatzer M. Patterns of care in metastatic pancreatic cancer: patient selection in clinical routine. *Therap Adv Gastroenterol* 2019; 12:1756284819877635.

Fitzal F, Gnant M, Greil R, Soelkner L, Schrenk P, Bartsch R, Singer C, Balic M, Hubalek M, Steger G, Knauer M, Bjelic-Radisic V, ABCSG. Impact of Breast Surgery in Primary Metastasized Breast Cancer: Outcomes of the Prospective Randomized Phase III ABCSG-28 POSYTTIVE Trial. *Ann Surg* 2018

Borchmann P, Bentz M, Keller U, Kühnhardt D, Ostermann H, Schmitz N, Hertenstein B, Aulitzky W, Maschmeyer G, Vieler T, Eich H, Baues C, Stein H, Fuchs M, Kuhnert G, Diehl V, Dietlein M, Hitz F, Topp M, Goergen H, Kobe C, Lohri A, Greil R, Eichenauer D, Zijlstra J, Markova J, Meissner J, Feuring-Buske M, Hüttmann A, Dierlamm J, Soekler M, Beck H, Willenbacher W, Ludwig W, Pabst T, Engert A. PET-guided treatment in patients with advanced-stage Hodgkin's lymphoma (HD18): final results of an open-label, international, randomised phase 3 trial by the German Hodgkin Study Group. *Lancet* 2017; 390:2790-2802.

Böll B, Engert A, Diehl V, Fuchs M, Borchmann S, Bürkle C, Eichenauer D, von Tresckow B, Greil R, Kerkhoff A, Hitz F, Bröckelmann P, Behringer K, Goergen H, Borchmann P. Bleomycin in older early-stage favorable Hodgkin lymphoma patients: analysis of the German Hodgkin Study Group (GHSg) HD10 and HD13 trials. *Blood* 2016; 127:2189-92.

Behringer K, Lange E, Mahlberg R, Hassler A, Vogelhuber M, Hahn D, Mezger J, Krause S, Skoetz N, Böll B, von Tresckow B, Diehl V, Hallek M, Borchmann P, Stein H, Eich H, Engert A, Bentz M, Vieler T, Kriz J, Goergen H, Tanner A, Zijlstra J, Greil R, Markova J, Sasse S, Fuchs M, Topp M, Soekler M, Mathas S, Meissner J, Wilhelm M, Koch P, Lindemann H, Schalk E, Semrau R. Omission of dacarbazine or bleomycin, or both, from the ABVD regimen in treatment of early-stage favourable Hodgkin's lymphoma (GHSg HD13): an open-label, randomised, non-inferiority trial. *Lancet* 2014; 385:1418-27.

Filipits M, Bauernhofer T, Hubalek M, Knauer M, Trapl H, Fesl C, Schaper C, Ferree S, Liu S, Cowens J, Gnant M, Dubsy P, Mlineritsch B, Singer C, Nielsen T, Rudas M, Greil R, Stöger H, Jakesz R, Bago-Horvath Z, Dietze O, Regitnig P, Gruber-Rossipal C, Müller-Holzner E, Austrian Breast and Colorectal Cancer Study Group. The PAM50 risk-of-recurrence score predicts risk for late distant recurrence after endocrine therapy in postmenopausal women with endocrine-responsive early breast cancer. *Clin Cancer Res* 2014; 20:1298-305.

Borchmann P, Paulus U, Smardova L, Huber C, Dörken B, Nerl C, Krause S, Mueller R, Fuchs M, Rank A, Greil R, Haverkamp H, Diehl V, Cerny T, Markova J, Ho A, Eich H, Mueller-Hermelink H, Kanz L, Engert A. Eight cycles of escalated-dose BEACOPP compared with four cycles of escalated-dose BEACOPP followed by four cycles of baseline-dose BEACOPP with or without radiotherapy in patients with advanced-stage Hodgkin's lymphoma: final analysis of the HD12 trial of the German Hodgkin Study Group. *J Clin Oncol* 2011; 29:4234-42.

Hofer S, Wick W, Zander T, Weder P, Roelcke U, Pichler J, Ochsenbein A, Marosi C, Lemke D, Huber U, Hottinger A, Greil R, Elandt K, Weller M. Clinical outcome with bevacizumab in patients with recurrent high-grade glioma treated outside clinical trials. *Acta Oncol* 2011; 50:630-5.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch