



Martin F Fey

Contact

Martin F Fey

Publications (9)

Kunstfeld R, Moser E, Mountzios G, Moynihan T, Nielsen T, Ohe Y, Öberg K, Palumbo A, Peccatori F, Pfeilstöcker M, Raut C, Michielin O, Meran J, Lenz H, Lichtman S, Licitra L, Lion T, Litière S, Liu L, Loehrer P, Markham M, Markman B, Mayerhoefer M, Remick S, Robson M, Rutkowski P, Stupp R, Sullivan R, Tabernero J, Travado L, Verheij M, Voest E, Vokes E, Von Roenn J, Weber J, Wildiers H, Strasser F, Stiefel F, Salgado R, Schapira L, Schernhammer E, Schlumberger M, Schmoll H, Schnipper L, Sessa C, Shapiro C, Steele J, Sternberg C, Yarden Y, Dittrich C, Bhatia S, Bogaerts J, Buckner J, Cardoso F, Casali P, Chu E, Close J, Coiffier B, Connolly R, Coupland S, Banerjee S, Awada A, Kosty M, Jezdic S, Pyle D, Berardi R, Bergh J, El-Saghir N, Lotz J, Österlund P, Pavlidis N, Purkalne G, De Petris L, De Santis M, de Vries E, Hahn S, Hanna D, Herold C, Herrstedt J, Homicsko K, Jones D, Jost L, Keilholz U, Khan S, Kiss A, Grothey A, Gopalan P, Dizon D, Duff J, Duska L, Eniu A, Ernstoff M, Felip E, Fey M, Gilbert J, Girard N, Glaudemans A, Köhne C. ESMO / ASCO Recommendations for a Global Curriculum in Medical Oncology Edition 2016. ESMO Open 2016; 1:e000097.

Pagani O, Goldhirsch A, Coates A, Fey M, Collins J, Thürlimann B, Lindtner J, Holmberg S, Castiglione-Gertsch M, Gelber R, Price K, International Breast Cancer Study Group (IBCSG). Patterns of recurrence of early breast cancer according to estrogen receptor status: a therapeutic target for a quarter of a century. Breast Cancer Res Treat 2009; 117:319-24.

Pagani O, Goldhirsch A, Coates A, Forbes J, Murray E, Fey M, Thürlimann B, Lindtner J, Collins J, Crivellari D, Holmberg S, Gelber R, Price K, Castiglione-Gertsch M, Simoncini E, Gelber S, International Breast Cancer Study Group. Is adjuvant chemotherapy of benefit for postmenopausal women who receive endocrine treatment for highly endocrine-responsive, node-positive breast cancer? International Breast Cancer Study Group Trials VII and 12-93. Breast Cancer Res Treat 2009; 116:491-500.

Karlsson P, Goldhirsch A, Gelber R, Forbes J, Crivellari D, Thürlimann B, Fey M, Holmberg S, Collins J, Lindtner J, Murray E, Gusterson B, Castiglione-Gertsch M, Coates A, Price K, Cole B, Wallgren A. The role of the number of uninvolved lymph nodes in predicting locoregional recurrence in breast cancer. JCO 2007; 25:2019-26.

Gruber G, Goldhirsch A, Coates A, Gelber R, Fey M, Simoncini E, Thürlimann B, Carbone A, Crivellari D, Collins J, Golouh R, Lindtner J, Holmberg S, Rudenstam C, Castiglione-Gertsch M, Price K, Nasi M, Bonetti M, International Breast Cancer Study Group. Prognostic value of extracapsular tumor spread for locoregional control in premenopausal patients with node-positive breast cancer treated with classical cyclophosphamide, methotrexate, and fluorouracil: long-term observations from International Breast Cancer Study Group Trial VI. JCO 2005; 23:7089-97.

Omlin A, Mühlemann K, Fey M, Pabst T. Pneumococcal vaccination in splenectomised cancer patients. European journal of cancer (Oxford, England : 1990) 2005; 41:1731-4.

Colleoni M, Thürlimann B, Crivellari D, Murray E, Mendiola C, Pagani O, Castiglione-Gertsch M, Coates A, Price K, Fey M, Collins J, Zahrieh D, Gelber R, Holmberg S, Mattsson J, Rudenstam C, Lindtner J, Erzen D, Snyder R, Goldhirsch A. Site of primary tumor has a prognostic role in operable breast cancer: the international breast cancer study group experience. JCO 2005; 23:1390-400.

Ghielmini M, Cerny T, Bargetzi M, Ketterer N, Stahel R, Pichert G, Schefer H, Betticher D, Fey M, Waltzer U, Bertoni F, Cogliatti S, Schmitz S, Swiss Group for Clinical Cancer Research. Effect of single-agent rituximab given at the standard schedule or as prolonged treatment in patients with mantle cell lymphoma: a study of the Swiss Group for Clinical Cancer Research (SAKK). Journal of clinical oncology : official journal of the American Society of Clinical Oncology 2005; 23:705-11.

Ghielmini M, Vorobiof D, Bargetzi M, Lohri A, Helg C, Kovacsovics T, Stupp R, Zucca E, Hess U, Peccatori F, Martinelli G, Betticher D, Fey M, Waltzer U, Hummerjohann J, Pichert G, Cogliatti S, Schmitz S, Cerny T. Prolonged treatment with rituximab in patients with follicular lymphoma significantly increases event-free survival and response duration compared with the standard weekly x 4 schedule. Blood 2004; 103:4416-23.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch