



Julia Dambacher

Contact

Julia Dambacher

Publications (15)

Seiderer J, Reinecker H, Lohse P, Ochsenkühn T, Göke B, Müller-Myhsok B, Jürgens M, Pfennig S, Niess J, Glas J, Tillack C, Leistner D, Dambacher J, Brand S. Genotype-phenotype analysis of the CXCL16 p.Ala181Val polymorphism in inflammatory bowel disease. *Clin Immunol* 2008; 127:49-55.

Dambacher J, Beigel F, Zitzmann K, Heeg M, Göke B, Diepolder H, Auernhammer C, Brand S. The role of interleukin-22 in hepatitis C virus infection. *Cytokine* 2008; 41:209-16.

Glas J, Klein W, Epplen J, Folwaczny C, Lohse P, Göke B, Ochsenkühn T, Mussack T, Folwaczny M, Müller-Myhsok B, Griga T, Haller D, Pfennig S, Konrad A, Schmechel S, Dambacher J, Seiderer J, Schroff F, Wetzke M, Roeske D, Török H, Tonenchi L, Brand S. The ATG16L1 gene variants rs2241879 and rs2241880 (T300A) are strongly associated with susceptibility to Crohn's disease in the German population. *Am J Gastroenterol* 2007; 103:682-91.

Glas J, Epplen J, Schiemann U, Folwaczny C, Lohse P, Göke B, Ochsenkühn T, Müller-Myhsok B, Folwaczny M, Mussack T, Klein W, Griga T, Maier K, Seiderer J, Wetzke M, Konrad A, Török H, Schmechel S, Tonenchi L, Grassl C, Dambacher J, Pfennig S, Brand S. rs1004819 is the main disease-associated IL23R variant in German Crohn's disease patients: combined analysis of IL23R, CARD15, and OCTN1/2 variants. *PloS one* 2007; 2:e819.

Dambacher J, Beigel F, Seiderer J, Haller D, Göke B, Auernhammer C, Brand S. Interleukin 31 mediates MAP kinase and STAT1/3 activation in intestinal epithelial cells and its expression is upregulated in inflammatory bowel disease. *Gut* 2007; 56:1257-65.

Brand S, Thasler W, Bilzer M, Diepolder H, Göke B, Storr M, Steib C, Olszak T, Prüfer T, Weiss T, Heeg M, Zitzmann K, Beigel F, Dambacher J, Auernhammer C. IL-22-mediated liver cell regeneration is abrogated by SOCS-1/3 overexpression in vitro. *Am J Physiol Gastrointest Liver Physiol* 2007; 292:G1019-28.

Dambacher J, Lohse P, Ochsenkühn T, Göke B, Diebold J, Otte J, Tillack C, Konrad A, Hofbauer K, Pfennig S, Schnitzler F, Sisic Z, Seiderer J, Staudinger T, Brand S. Macrophage migration inhibitory factor (MIF) -173G/C promoter polymorphism influences upper gastrointestinal tract involvement and disease activity in patients with Crohn's disease. *Inflamm Bowel Dis* 2007; 13:71-82.

Seiderer J, Lohse P, Göke B, Sackmann M, Tillack C, Dambacher J, Hofbauer K, Herrmann K, Pfennig S, Staudinger T, Brand S, Schnitzler F, Ochsenkühn T. Homozygosity for the CARD15 frameshift mutation 1007fs is predictive of early onset of Crohn's disease with ileal stenosis, entero-enteral fistulas, and frequent need for surgical intervention with high risk of re-stenosis. *Scand J Gastroenterol* 2006; 41:1421-32.

Brand S, Auernhammer C, Göke B, Ochsenkühn T, Seiderer J, Herrmann K, Leclair S, Popp A, Jagla W, Marquardt A, Diepolder H, Otte J, Eichhorst S, Zitzmann K, Olszak T, Beigel F, Dambacher J. IL-22 is increased in active Crohn's disease and promotes proinflammatory gene expression and intestinal epithelial cell migration. *Am J Physiol Gastrointest Liver Physiol* 2006; 290:G827-38.

Brand S, Olszak T, Beigel F, Diebold J, Otte J, Eichhorst S, Göke B, Dambacher J. Cell differentiation dependent expressed CCR6 mediates ERK-1/2, SAPK/JNK, and Akt signaling resulting in proliferation and migration of colorectal cancer cells. *J Cell Biochem* 2006; 97:709-23.

Brand S, Ochsenkühn T, Göke B, Konrad A, Tillack C, Seiderer J, Pfennig S, Staudinger T, Schnitzler F, Dambacher J, Hofbauer K, Lohse P. Increased expression of the chemokine fractalkine in Crohn's disease and association of the fractalkine receptor T280M polymorphism with a fibrostenosing disease Phenotype. *Am J Gastroenterol* 2006; 101:99-106.

Brand S, Dambacher J, Beigel F, Olszak T, Diebold J, Otte J, Göke B, Eichhorst S. CXCR4 and CXCL12 are inversely expressed in colorectal cancer cells and modulate cancer cell migration, invasion and MMP-9 activation. *Exp Cell Res* 2005; 310:117-30.

Brand S, Göke B, Auernhammer C, Adler B, Diepolder H, Diebold J, Otte J, Eichhorst S, Zitzmann K, Olszak T, Beigel F, Dambacher J. IL-28A and IL-29 mediate antiproliferative and antiviral signals in intestinal epithelial cells and murine CMV infection increases colonic IL-28A expression. *Am J Physiol Gastrointest Liver Physiol* 2005; 289:G960-8.

Brand S, Lohse P, Göke B, Crispin A, Konrad A, Tillack C, Seiderer J, Dambacher J, Hofbauer K, Pfennig S, Schnitzler F, Staudinger T, Ochsenkühn T. The role of Toll-like receptor 4 Asp299Gly and Thr399Ile polymorphisms and CARD15/NOD2 mutations in the susceptibility and phenotype of Crohn's disease. *Inflamm Bowel Dis* 2005; 11:645-52.

Brand S, Diepolder H, Göke B, Eichhorst S, Vlotides G, Olszak T, Beigel F, Dambacher J, Zitzmann K, Auernhammer C. SOCS-1 inhibits expression of the antiviral proteins 2',5'-OAS and MxA induced by the novel interferon-lambda IL-28A and IL-29. *Biochem Biophys Res Commun* 2005; 331:543-8.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch