



Maria Weidinger

Contact

Maria Weidinger

Publications (8)

Beigel F, Ochsenkühn T, Seiderer J, Göke B, Weidinger M, Breitenreicher S, Schnitzler F, Tillack C, Laubender R, Löhr B, Brand S. Iron status and analysis of efficacy and safety of ferric carboxymaltose treatment in patients with inflammatory bowel disease. *Digestion* 2011; 85:47–54.

Glas J, Czamara D, Diegelmann J, Müller-Myhsok B, Lohse P, Wolf C, Ochsenkühn T, Göke B, Lass U, Olszak T, Beigel F, Weidinger M, Pfennig S, Tillack C, Fries C, Seiderer J, Brand S. CEACAM6 gene variants in inflammatory bowel disease. *PloS one* 2011; 6:e19319.

Beigel F, Schnitzler F, Paul Laubender R, Pfennig S, Weidinger M, Göke B, Seiderer J, Ochsenkühn T, Brand S. Formation of antinuclear and double-strand DNA antibodies and frequency of lupus-like syndrome in anti-TNF- α antibody-treated patients with inflammatory bowel disease. *Inflamm Bowel Dis* 2011; 17:91–8.

Glas J, Czamara D, Diegelmann J, Lohse P, Ochsenkühn T, Göke B, Müller-Myhsok B, Weidinger M, Laubender R, Olszak T, Jürgens M, Beigel F, Pfennig S, Tillack C, Seiderer J, Brand S. The NOD2 single nucleotide polymorphisms rs2066843 and rs2076756 are novel and common Crohn's disease susceptibility gene variants. *PloS one* 2010; 5:e14466.

Glas J, Czamara D, Diegelmann J, Müller-Myhsok B, Folwaczny M, Ochsenkühn T, Göke B, Weidinger M, Olszak T, Beigel F, Wetzke M, Pfennig S, Tengler B, Fischer D, Seiderer J, Brand S. Pregnane X receptor (PXR/NR112) gene haplotypes modulate susceptibility to inflammatory bowel disease. *Inflamm Bowel Dis* 2010; 17:1917–24.

Glas J, Roeske D, Müller-Myhsok B, Diegelmann J, Ochsenkühn T, Göke B, Folwaczny M, Lohse P, Epplen J, Klein W, Pfennig S, Weidinger M, Beigel F, Fries C, Nagy M, Seiderer J, Brand S. Evidence for STAT4 as a common autoimmune gene: rs7574865 is associated with colonic Crohn's disease and early disease onset. *PloS one* 2010; 5:e10373.

Jürgens M, Herrmann K, Lohse P, Göke B, Kreis M, Schnitzler F, Weidinger M, Beigel F, Tillack C, Pfennig S, Wagner J, Wetzke M, Glas J, Seiderer J, Laubender R, Brand S, Ochsenkühn T. The presence of fistulas and NOD2 homozygosity strongly predict intestinal stenosis in Crohn's disease independent of the IL23R genotype. *J Gastroenterol* 2010; 45:721–31.

Jürgens M, Ochsenkühn T, Glas J, Göke B, Lohse P, Tillack C, Schnitzler F, Stallhofer J, Pfennig S, Beigel F, Wetzke M, Wagner J, Seiderer J, Weidinger M, Hartl F, Laubender R, Brand S. Disease activity, ANCA, and IL23R genotype status determine early response to infliximab in patients with ulcerative colitis. *Am J Gastroenterol* 2010; 105:1811–9.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch