



Johanna Wagner

Contact

Johanna Wagner

Publications (5)

Tillack C, Koburger M, Wagner J, Glas J, Diegelmann J, Koglin S, Dombrowski Y, Schaubert J, Wollenberg A, Maier H, Wetzke M, Ehmann L, Friedrich M, Laubender R, Papay P, Vogelsang H, Stallhofer J, Beigel F, Bedynek A, Brand S. Anti-TNF antibody-induced psoriasiform skin lesions in patients with inflammatory bowel disease are characterised by interferon- γ -expressing Th1 cells and IL-17A/IL-22-expressing Th17 cells and respond to anti-IL-12/IL-23 antibody treatment. *Gut* 2013; 63:567-77.

Glas J, Czamara D, Diegelmann J, Ochsenkühn T, Göke B, Wetzke M, Steib C, Stallhofer J, Beigel F, Friedrich M, Tillack C, Fries C, Olszak T, Wagner J, Seiderer J, Brand S. Analysis of IL12B gene variants in inflammatory bowel disease. *PloS one* 2012; 7:e34349.

Glas J, Czamara D, Diegelmann J, Karbalai N, Ochsenkühn T, Göke B, Steib C, Friedrich M, Stallhofer J, Tillack C, Beigel F, Wetzke M, Olszak T, Seiderer J, Wagner J, Brand S. PTPN2 gene variants are associated with susceptibility to both Crohn's disease and ulcerative colitis supporting a common genetic disease background. *PloS one* 2012; 7:e33682.

Jürgens M, Herrmann K, Lohse P, Göke B, Kreis M, Schnitzler F, Weidinger M, Beigel F, Tillack C, Pfennig S, Wagner J, Wetzke M, Glas J, Seiderer J, Laubender R, Brand S, Ochsenkühn T. The presence of fistulas and NOD2 homozygosity strongly predict intestinal stenosis in Crohn's disease independent of the IL23R genotype. *J Gastroenterol* 2010; 45:721-31.

Jürgens M, Ochsenkühn T, Glas J, Göke B, Lohse P, Tillack C, Schnitzler F, Stallhofer J, Pfennig S, Beigel F, Wetzke M, Wagner J, Seiderer J, Weidinger M, Hartl F, Laubender R, Brand S. Disease activity, ANCA, and IL23R genotype status determine early response to infliximab in patients with ulcerative colitis. *Am J Gastroenterol* 2010; 105:1811-9.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch