



**L Philip Schumm**

**Contact**

L Philip Schumm

## Publications (4)

Akhlaghpour M, Haritunians T, More S, Thomas L, Stamps D, Dube S, Li D, Yang S, Landers C, Mengesha E, Hamade H, Murali R, Potdar A, Wolf A, Botwin G, Khrom M, International IBD Genetics Consortium, Ananthakrishnan A, Faubion W, Jabri B, Lira S, Newberry R, Sandler R, Sartor R, Xavier R, Brant S, Cho J, Duerr R, Lazarev M, Rioux J, Schumm L, Silverberg M, Zaghiyan K, Fleshner P, Melmed G, Vasiliauskas E, Ha C, Rabizadeh S, Syal G, Bonthala N, Ziring D, Targan S, Long M, McGovern D, Michelsen K, Brand S. Genetic coding variant in complement factor B (CFB) is associated with increased risk for perianal Crohn's disease and leads to impaired CFB cleavage and phagocytosis. *Gut* 2023; 72:2068-2080.

Li D, Devlin B, Sharma Y, Torkvist L, Targan S, Stempak J, Simms L, Regueiro M, Proctor D, Borneman J, Hakonarson H, McGovern D, Braun J, Cho J, Silverberg M, Rioux J, Brant S, Daly M, Xavier R, Milgrom R, Glas J, Halfvarson J, Radford-Smith G, Brand S, D'Amato M, Hui K, Jacobs J, Haritunians T, Achkar J, Niess J, Kugathasan S, Fiocchi C, Dubinsky M, Baidoo L, Aumais G, Ananthakrishnan A, Klei L, Schumm L, Büning C, Duerr R. A Pleiotropic Missense Variant in SLC39A8 Is Associated With Crohn's Disease and Human Gut Microbiome Composition. *Gastroenterology* 2016; 151:724-32.

Lees C, van der Meulen-de Jong A, Theatre E, Schreiber S, Satsangi J, Lee J, Lawrance I, Kupcinkas L, Kennedy N, Weersma R, Wilson D, Barrett J, McGovern D, Radford-Smith G, Silverberg M, Mansfield J, Rioux J, Vermeire S, Parkes M, Huang H, Hov J, Halfvarson J, Annese V, Andrews J, Andersen V, Ahmad T, Zeissig S, Schumm L, Jostins L, Boucher G, Brand S, Brant S, Cho J, Hakonarson H, Goyette P, Geary R, Franke A, Ferguson L, Duerr R, Dubinsky M, Daly M, Cleyen I. Inherited determinants of Crohn's disease and ulcerative colitis phenotypes: a genetic association study. *Lancet* 2015; 387:156-67.

Kupcinkas L, Potocnik U, Prescott N, Regueiro M, Rotter J, Russell R, Sanderson J, Sans M, Satsangi J, Schreiber S, Simms L, Sventoraityte J, Ponsioen C, Palmieri O, Kugathasan S, Latiano A, Laukens D, Lawrance I, Lees C, Louis E, Mahy G, Mansfield J, Morgan A, Mowat C, Newman W, Targan S, Taylor K, Tremelling M, Hakonarson H, Brant S, Radford-Smith G, Mathew C, Rioux J, Schadt E, Daly M, Franke A, Parkes M, Vermeire S, Barrett J, Annese V, Silverberg M, Verspaget H, De Vos M, Wijmenga C, Wilson D, Winkelmann J, Xavier R, Zeissig S, Zhang B, Zhang C, Zhao H, Cho J, Karlsen T, Jostins L, Theatre E, Spain S, Raychaudhuri S, Goyette P, Wei Z, Abraham C, Achkar J, Ahmad T, Amininejad L, Ananthakrishnan A, Andersen V, Cleyen I, Ning K, Ripke S, Weersma R, Duerr R, McGovern D, Hui K, Lee J, Schumm L, Sharma Y, Anderson C, Essers J, Mitrovic M, Andrews J, Baidoo L, Balschun T, Ferguson L, Franchimont D, Fransen K, Geary R, Georges M, Gieger C, Glas J, Haritunians T, Hart A, Hawkey C, Hedl M, Ellinghaus D, Edwards C, Bampton P, Bitton A, Boucher G, Brand S, Büning C, Cohain A, Cichon S, D'Amato M, De Jong D, Devaney K, Dubinsky M, Hu X. Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 2012; 491:119-24.

## Projects (0)

No results found.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)