



Max M Lenders

Contact

Max M Lenders

Publications (4)

Wagner N, Lenders M, Kühl K, Reinhardt L, Fuchß M, Ring N, Stäger R, Zellweger C, Ebel C, Kimeswenger S, Oellinger A, Amaral T, Forschner A, Leiter U, Klumpp B, Hoetzenecker W, Terheyden P, Mangana J, Loquai C, Cozzio A, Garbe C, Meier F, Eigentler T, Flatz L. Baseline metastatic growth rate is an independent prognostic marker in patients with advanced BRAF V600 mutated melanoma receiving targeted therapy. *Eur J Cancer* 2023; 196:113425.

Gassenmaier M, Lenders M, Forschner A, Leiter U, Weide B, Garbe C, Eigentler T, Wagner N. Serum S100B and LDH at Baseline and During Therapy Predict the Outcome of Metastatic Melanoma Patients Treated with BRAF Inhibitors. *Target Oncol* 2021; 16:197-205.

Wagner N, Schraag A, Klumpp B, Hoetzenecker W, Berking C, Richtig E, Ziemer M, Mangana J, Terheyden P, Loquai C, Nguyen V, Gebhardt C, Meier F, Diem S, Cozzio A, Flatz L, Röcken M, Garbe C, Gassenmaier M, Weide B, Leiter U, Lenders M, Kühl K, Reinhardt L, André F, Dudda M, Ring N, Ebel C, Stäger R, Zellweger C, Lang R, Paar M, Gussek P, Richtig G, Stürmer S, Kimeswenger S, Oellinger A, Forschner A, Eigentler T. Pretreatment metastatic growth rate determines clinical outcome of advanced melanoma patients treated with anti-PD-1 antibodies: a multicenter cohort study. *J Immunother Cancer* 2021; 9

Winter J, Eigentler T, Garbe C, Röcken M, Cozzio A, Flatz L, Purde M, Weide B, Leiter U, Forschner A, Gassenmaier M, Lenders M, Wagner N. Prognostic role of gamma-glutamyl transferase in metastatic melanoma patients treated with immune checkpoint inhibitors. *Cancer Immunol Immunother* 2020; 70:1089-1099.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch