



C Hanusch

Contact

C Hanusch

Publications (9)

Loibl S, Staib P, Link T, Rhiem K, Solbach C, Fasching P, Nekljudova V, Denkert C, Untch M, Jackisch C, Thomalla J, Schneeweiss A, Huober J, Braun M, Rey J, Blohmer J, Furlanetto J, Zahm D, Hanusch C, GBG and AGO-B. Neoadjuvant durvalumab improves survival in early triple-negative breast cancer independent of pathological complete response. *Ann Oncol* 2022;

Loibl S, Schmitt W, von Minckwitz G, Thomalla J, Kümmel S, Rautenberg B, Fasching P, Weber K, Rhiem K, Denkert C, Jackisch C, Rezai M, Engels K, Untch M, Burchardi N, Huober J, Sinn B, Blohmer J, Grischke E, Furlanetto J, Tesch H, Hanusch C, Schneeweiss A. Corrigendum to "A randomised phase II study investigating durvalumab in addition to an anthracycline taxane-based neoadjuvant therapy in early triple-negative breast cancer: clinical results and biomarker analysis of GeparNuevo study": [Annals of Oncology (2019), volume 30:1279-1288]. *Ann Oncol* 2022; 33:743-744.

Fasching P, Rhiem K, Schneeweiss A, Engels K, Denkert C, Schmutzler R, Hahnen E, Untch M, Burchardi N, Blohmer J, Loibl S, Doering G, Uleer C, Schmitt W, Link T, Hauke J, Seither F, Jackisch C, Klare P, Schmatloch S, Hanusch C, Huober J, Stefek A, Seiler S, German Breast Group and Arbeitsgemeinschaft Gynäkologische Onkologie Breast. Neoadjuvant paclitaxel/olaparib in comparison to paclitaxel/carboplatinum in patients with HER2-negative breast cancer and homologous recombination deficiency (GeparOLA study). *Ann Oncol* 2020; 32:49-57.

Karn T, Schmitt W, Wu S, van Mackelenbergh M, Müller V, Schem C, Stickeler E, Fasching P, Jackisch C, Untch M, Schneeweiss A, Marmé F, Blohmer J, Denkert C, Weber K, Holtrich U, Hanusch C, Sinn B, Higgs B, Jank P, Sinn H, Huober J, Becker C, Loibl S. Tumor mutational burden and immune infiltration as independent predictors of response to neoadjuvant immune checkpoint inhibition in early TNBC in GeparNuevo. *Ann Oncol* 2020; 31:1216-1222.

Loibl S, Schmitt W, von Minckwitz G, Thomalla J, Kümmel S, Rautenberg B, Fasching P, Weber K, Rhiem K, Denkert C, Jackisch C, Rezai M, Engels K, Untch M, Burchardi N, Huober J, Sinn B, Blohmer J, Grischke E, Furlanetto J, Tesch H, Hanusch C, Schneeweiss A. A randomised phase II study investigating durvalumab in addition to an anthracycline taxane-based neoadjuvant therapy in early triple-negative breast cancer: clinical results and biomarker analysis of GeparNuevo study. *Ann Oncol* 2019; 30:1279-1288.

von Minckwitz G, Jackisch C, Kümmel S, Fasching P, Schneeweiss A, Paepke S, Untch M, Burchardi N, Mehta K, Loibl S, Hanusch C, Denkert C, Rezai M, Tesch H, Huober J, Gerber B, Zahm D, Hilfrich J, Costa S, Dubsky P, Blohmer J, German Breast Group and Austrian Breast and Colon Cancer Study Group Investigators. Zoledronate for patients with invasive residual disease after anthracyclines-taxane-based chemotherapy for early breast cancer - The Phase III NeoAdjuvant Trial Add-oN (NaTaN) study (GBG 36/ABCSCG 29). *Eur J Cancer* 2016; 64:12-21.

Deuker J, Kohls A, Weikel W, Bischoff J, Freese K, Schmidt M, Wiest W, Sütterlin M, Dietrich M, Grießhammer M, Burgmann D, Rack B, Salat C, Sattler D, Tio J, von Abel E, Fischer D, Ober A, Strumberg D, Uleer C, Solomayer E, Runnebaum I, Link H, Tomé O, Ulmer H, Conrad B, Feisel-Schwickardi G, Schumacher C, Steinmetz T, Bauerfeind I, Kremers S, Langanke D, Kullmer U, Christensen B, Burkamp U, Hofmann M, Schuster J, Simon W, Kronawitter U, Clemens M, Janni W, Latos K, Bauer W, Roßmann A, Bauer L, Lampe D, Heyl V, Hoffmann G, Lorenz-Salehi F, Hackmann J, Ruhland F, Stefek A, Köhne C, Meinerz W, Grasshoff S, Decker T, Overkamp F, Thalmann I, Sallmann A, Beck T, Reimer T, Bartzke G, Deryal M, Weigel M, Weder P, Steffens C, Lemster S, Schlag R, Müller T, von Minckwitz G, Nekljudova V, Gerber B, GBG/AGO-B study groups, Gnauert K, Heinrich B, Prätz T, Groh U, Tanzer H, Villena C, Tulusan A, Liedtke B, Blohmer J, Mau C, Potenberg J, Schilling J, Fehm T, Hauschild M, Loibl S, Untch M, Eidtmann H, Rezai M, Fasching P, Tesch H, Eggemann H, Schrader I, Kittel K, Hanusch C, Huober J, Solbach C, Jackisch C, Kunz G, Blohmer J, Just M, Weiss E, Möbus V, Stickeler E, Heinrich G, Wagner H, Abdallah A, Dewitz T, Emons G, Belau A, Rethwisch V, Lantzsch T, Thomssen C, Mattner U, Nugent A, Müller V, Noesselt T, Khandan F, Kühn T, Bückner U, Wolfgarten M, Lorenz R, Doering G, Feidicker S, Krabisch P, Deichert U, Augustin D, Kast K, Nestle-Krämling C, Höß C, Terhaag J, Fasching P, Staib P, Aktas B, Holms F. Survival after neoadjuvant chemotherapy with or without bevacizumab or everolimus for HER2-negative primary breast cancer (GBG 44-GeparQuinto)†. *Ann Oncol* 2014; 25:2363-2372.

von Minckwitz G, Costa S, Jackisch C, Paepke S, Schneeweiss A, Kümmel S, Denkert C, Mehta K, Loibl S, Blohmer J, Hanusch C, Rezai M, Fasching P, Huober J, Tesch H, Bauerfeind I, Hilfrich J, Eidtmann H, Gerber B, Untch M. Survival after adding capecitabine and trastuzumab to neoadjuvant anthracycline-taxane-based chemotherapy for primary breast cancer (GBG 40--GeparQuattro). *Ann Oncol* 2013; 25:81-9.

Gerber B, Jackisch C, Kunz G, Blohmer J, Huober J, Hauschild M, Nekljudova V, Untch M, von Minckwitz G, Solbach C, Kreienberg R, Hanusch C, Loibl S, Eidtmann H, Rezai M, Fasching P, Tesch H, Eggemann H, Schrader I, Kittel K, German Breast Group Investigators. Neoadjuvant bevacizumab and anthracycline-taxane-based chemotherapy in 678 triple-negative primary breast cancers; results from the geparquinto study (GBG 44). *Ann Oncol* 2013; 24:2978-84.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch