



Stefan Paepke

Contact

Stefan Paepke

Publications (14)

Pfob A, Cai L, Schneeweiss A, Rauch G, Thomas B, Schaeffgen B, Kümmel S, Reimer T, Hahn M, Thill M, Blohmer J, Hackmann J, Malter W, Bekes I, Friedrichs K, Wojcinski S, Joos S, Paepke S, Degenhardt T, Rom J, Rody A, van Mackelenbergh M, Banys-Paluchowski M, Große R, Reinisch M, Karsten M, Sidey-Gibbons C, Wallwiener M, Golatta M, Heil J. ASO Visual Abstract: Minimally-Invasive Breast Biopsy After Neoadjuvant Systemic Treatment to Identify Breast Cancer Patients with Residual Disease for Extended Neoadjuvant Treatment-A New Concept. *Ann Surg Oncol* 2023

Pfob A, Cai L, Schneeweiss A, Rauch G, Thomas B, Schaeffgen B, Kümmel S, Reimer T, Hahn M, Thill M, Blohmer J, Hackmann J, Malter W, Bekes I, Friedrichs K, Wojcinski S, Joos S, Paepke S, Degenhardt T, Rom J, Rody A, van Mackelenbergh M, Banys-Paluchowski M, Große R, Reinisch M, Karsten M, Sidey-Gibbons C, Wallwiener M, Golatta M, Heil J. Minimally Invasive Breast Biopsy After Neoadjuvant Systemic Treatment to Identify Breast Cancer Patients with Residual Disease for Extended Neoadjuvant Treatment: A New Concept. *Ann Surg Oncol* 2023

Pfob A, Joos S, Paepke S, Degenhardt T, Rom J, Rody A, van Mackelenbergh M, Banys-Paluchowski M, Große R, Reinisch M, Karsten M, Golatta M, Wojcinski S, Friedrichs K, Bekes I, Sidey-Gibbons C, Rauch G, Thomas B, Schaeffgen B, Kuemmel S, Reimer T, Hahn M, Thill M, Blohmer J, Hackmann J, Malter W, Heil J. Intelligent Vacuum-Assisted Biopsy to Identify Breast Cancer Patients With Pathologic Complete Response (ypT0 and ypN0) After Neoadjuvant Systemic Treatment for Omission of Breast and Axillary Surgery. *J Clin Oncol* 2022;JCO2102439.

Heil J, Friedrichs K, Wojcinski S, Joos S, Paepke S, Ditsch N, Rody A, Große R, van Mackelenbergh M, Reinisch M, Karsten M, Golatta M, Bekes I, Malter W, Hackmann J, Pfob A, Sinn H, Rauch G, Bach P, Thomas B, Schaeffgen B, Kuemmel S, Reimer T, Hahn M, Thill M, Blohmer J, RESPONDER Investigators. Diagnosing Pathologic Complete Response in the Breast After Neoadjuvant Systemic Treatment of Breast Cancer Patients by Minimal Invasive Biopsy: Oral Presentation at the San Antonio Breast Cancer Symposium on Friday, December 13, 2019, Program Number GS5-03. *Ann Surg* 2020

Krug D, Huober J, Jackisch C, Kümmel S, Paepke S, Schem C, Schneeweiss A, Untch M, Debus J, von Minckwitz G, Kühn T, Hilfrich J, Heil J, Lederer B, Seither F, Nekljudova V, Ataseven B, Blohmer J, Costa S, Denkert C, Ditsch N, Gerber B, Hanusch C, Loibl S. Post-Mastectomy Radiotherapy After Neoadjuvant Chemotherapy in Breast Cancer: A Pooled Retrospective Analysis of Three Prospective Randomized Trials. *Ann Surg Oncol* 2019; 26:3892-3901.

Hahnen E, Huober J, Rhiem K, Costa S, Altmüller J, Hanusch C, Thiele H, Müller V, Nürnberg P, Karn T, Nekljudova V, Untch M, von Minckwitz G, Neidhardt G, Schem C, Lederer B, Hauke J, Loibl S, Kröber S, Schneeweiss A, Denkert C, Fasching P, Blohmer J, Jackisch C, Paepke S, Gerber B, Kümmel S, Schmutzler R. Germline Mutation Status, Pathological Complete Response, and Disease-Free Survival in Triple-Negative Breast Cancer: Secondary Analysis of the GeparSixto Randomized Clinical Trial. *JAMA Oncol* 2017; 3:1378-1385.

Kümmel S, Burchardi N, von Minckwitz G, Fasching P, Klare P, Engels K, Denkert C, Schneeweiss A, Jackisch C, Hilfrich J, Hanusch C, Gerber B, Eiermann W, Blohmer J, Untch M, Schem C, Huober J, Paepke S, Loibl S. Randomised, open-label, phase II study comparing the efficacy and the safety of cabazitaxel versus weekly paclitaxel given as neoadjuvant treatment in patients with operable triple-negative or luminal B/HER2-negative breast cancer (GENEVIEVE). *Eur J Cancer* 2017; 84:1-8.

Marme F, von Minckwitz G, Untch M, Paepke S, Loibl S, Kümmel S, Jackisch C, Huober J, Hilfrich J, Hanusch C, Gerber B, Eidtmann H, Denkert C, Costa S, Blohmer J, Lederer B, Schneeweiss A. Utility of the CPS+EG staging system in hormone receptor-positive, human epidermal growth factor receptor 2-negative breast cancer treated with neoadjuvant chemotherapy. *Eur J Cancer* 2015; 53:65-74.

Loibl S, Mehta K, Nekljudova V, Blohmer J, Costa S, Denkert C, Eidtmann H, Gerber B, Hanusch C, Hilfrich J, Huober J, Schneeweiss A, Kümmel S, Paepke S, Untch M, Lederer B, Jackisch C, von Minckwitz G. Outcome after neoadjuvant chemotherapy in young breast cancer patients: a pooled analysis of individual patient data from eight prospectively randomized controlled trials. *Breast Cancer Res Treat* 2015; 152:377-87.

Hanusch C, Burchardi N, Engels K, Blohmer J, Costa S, Denkert C, Eidtmann H, Gerber B, Hilfrich J, Huober J, Jackisch C, Kümmel S, Paepke S, Untch M, Loibl S, Schneeweiss A, von Minckwitz G. Dual Blockade with Afatinib and Trastuzumab as Neoadjuvant Treatment for Patients with Locally Advanced or Operable Breast Cancer Receiving Taxane-Anthracycline Containing Chemotherapy-DAFNE (GBG-70). *Clin Cancer Res* 2015; 21:2924-31.

Fontanella C, Schneeweiss A, Paepke S, Jackisch C, Mehta K, Nekljudova V, Untch M, Neven P, von Minckwitz G, Huober J, Hilfrich J, Hanusch C, Lederer B, Gade S, Vanoppen M, Blohmer J, Costa S, Denkert C, Eidtmann H, Gerber B, Loibl S. Impact of body mass index on neoadjuvant treatment outcome: a pooled analysis of eight prospective neoadjuvant breast cancer trials. *Breast Cancer Res Treat* 2015; 150:127-39.

Loibl S, Fasching P, Andre F, Lindner J, Sotiriou C, Dykgers A, Guo S, Gade S, Nekljudova V, Loi S, Untch M, Pfitzner B, Huober J, Heinrichs C, von Minckwitz G, Schneeweiss A, Paepke S, Lehmann A, Rezai M, Zahm D, Sinn P, Khandan F, Eidtmann H, Dohnal K, Denkert C. PIK3CA mutations are associated with lower rates of pathologic complete response to anti-human epidermal growth factor receptor 2 (her2) therapy in primary HER2-overexpressing breast cancer. *J Clin Oncol* 2014; 32:3212-20.

von Minckwitz G, Huober J, Costa S, Tesch H, Hanusch C, Hilfrich J, Khandan F, Fasching P, Sinn B, Engels K, Mehta K, Nekljudova V, Klare P, Eidtmann H, Schneeweiss A, Loibl S, Salat C, Denkert C, Rezai M, Blohmer J, Jackisch C, Paepke S, Gerber B, Zahm D, Kümmel S, Untch M. Neoadjuvant carboplatin in patients with triple-negative and HER2-positive early breast cancer (GeparSixto; GBG 66): a randomised phase 2 trial. *Lancet Oncol* 2014; 15:747-56.

von Minckwitz G, Mehta K, Zahm D, Untch M, Schneeweiss A, Paepke S, Kümmel S, Kaufmann M, Jackisch C, Huober J, Hilfrich J, Hanusch C, Gerber B, Eiermann W, Eidtmann H, Denkert C, Costa S, Blohmer J, Loibl S. Response-guided neoadjuvant chemotherapy for breast cancer. *J Clin Oncol* 2013; 31:3623-30.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch