



**D Weishaupt**

**Contact**

D Weishaupt

## Publications (9)

Donati O, Reiner C, Hany T, Fornaro J, von Schulthess G, Marincek B, Weishaupt D. 18F-FDG-PET and MRI in patients with malignancies of the liver and pancreas. Accuracy of retrospective multimodality image registration by using the CT-component of PET/CT. *Nuklearmedizin* 2010; 49:106-14.

Hausammann R, Steffen T, Weishaupt D, Beutner U, Hetzer F. Rectocele and intussusception: is there any coherence in symptoms or additional pelvic floor disorders?. *Techniques in coloproctology* 2009; 13:17-25; discussion 25-6.

Hübner M, Hetzer F, Weishaupt D, Hahnloser D, Clavien P, Demartines N. A prospective comparison between clinical outcome and open-configuration magnetic resonance defecography findings before and after surgery for symptomatic rectocele. *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland* 2006; 8:605-11.

Roos J, Desbiolles L, Weishaupt D, Wildermuth S, Hilfiker P, Marincek B, Boehm T. Multi-detector row CT: effect of iodine dose reduction on hepatic and vascular enhancement. *Rofo* 2004; 176:556-63.

Zanetti M, Carstensen T, Weishaupt D, Jost B, Hodler J. MR arthrographic variability of the arthroscopically normal glenoid labrum: qualitative and quantitative assessment. *Eur Radiol* 2001; 11:559-66.

Weishaupt D, Hetzer F, Ruehm S, Patak M, Schmidt M, Debatin J. Three-dimensional contrast-enhanced MRI using an intravascular contrast agent for detection of traumatic intra-abdominal hemorrhage and abdominal parenchymal injuries: an experimental study. *European radiology* 2000; 10:1958-64.

Hilfiker P, Weishaupt D, Kacel G, Hetzer F, Griff M, Ruehm S, Debatin J. Comparison of three dimensional magnetic resonance imaging in conjunction with a blood pool contrast agent and nuclear scintigraphy for the detection of experimentally induced gastrointestinal bleeding. *Gut* 1999; 45:581-7.

Weishaupt D, Wildermuth S, Schmid M, Hilfiker P, Hodler J, Debatin J. Virtual MR arthroscopy: new insights into joint morphology. *Journal of magnetic resonance imaging : JMRI* 1999; 9:757-60.

Zanetti M, Weishaupt D, Jost B, Gerber C, Hodler J. MR imaging for traumatic tears of the rotator cuff: high prevalence of greater tuberosity fractures and subscapularis tendon tears. *AJR Am J Roentgenol* 1999; 172:463-7.

## Projects (0)

No results found.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)