



Wolfgang Wick

Contact

Wolfgang Wick

Publications (11)

German Consortium on Aggressive Meningiomas (KAM), Ratliff M, Neidert M, Wirsching H, Harter P, Unterberg A, Herold-Mende C, Jungwirth G, Etminan N, Lamszus K, Westphal M, Hänggi D, Jungk C, Mawrin C, Platten M, Snuderl M, Sahm F, von Deimling A, Preusser M, Weller M, Wick W, Acker T, Reifenberger G, Jones D, Pfister S, Frank S, Hench J, Brandner S, Aldape K, Sen C, Golfinos J, Serrano J, Stein M, Dohmen H, Reuss D, Dogan H, Patel A, Blume C, Euskirchen P, Sill M, Schrimpf D, Berghoff A, Sievers P, Hielscher T, Stichel D, Reinhardt A, Suwala A, Grady C, Jones T, Bridges L, Greenway F, Leu S, Jaunmuktane Z, Schittenhelm J, Ketter R, Bewerunge-Hudler M, Rushing E, Ricklefs F, Baumgarten P, Wefers A, Maas S. Integrated Molecular-Morphologic Meningioma Classification: A Multicenter Retrospective Analysis, Retrospectively and Prospectively Validated. *J Clin Oncol* 2021; 39:3839–3852.

Sievers P, Etminan N, Ratliff M, Herold-Mende C, Pfister S, Jones D, Weller M, Harter P, Wick W, Preusser M, von Deimling A, Westphal M, Lamszus K, Reifenberger G, Hielscher T, Schrimpf D, Stichel D, Reuss D, Berghoff A, Neidert M, Wirsching H, Mawrin C, Ketter R, Paulus W, Sahm F. CDKN2A/B homozygous deletion is associated with early recurrence in meningiomas. *Acta Neuropathol* 2020; 140:409–413.

van den Bent M, Zhang H, Balbin O, Vicente S, Cheng Y, Kirsilae T, Zhao S, Tiedt R, Rodón J, Tabatabai G, Jörger M, Lassman A, Wen P, Yung W, Sepulveda J, De Vos F, Azaro A, Wick W. A Phase Ib/II, open-label, multicenter study of INC280 (capmatinib) alone and in combination with buparlisib (BKM120) in adult patients with recurrent glioblastoma. *J Neurooncol* 2019

Paramasivam N, Beck K, Brehmer S, Urbschat S, Seiz-Rosenhagen M, Hänggi D, Herold-Mende C, Ketter R, Eils R, Ram Z, Pfister S, Wick W, Weller M, Grossmann R, von Deimling A, Schlesner M, Bewerunge-Hudler M, Schick M, Hübschmann D, Toprak U, Ishaque N, Neidert M, Schrimpf D, Stichel D, Reuss D, Sievers P, Reinhardt A, Wefers A, Jones D, Gu Z, Werner J, Uhrig S, Wirsching H, Sahm F. Mutational patterns and regulatory networks in epigenetic subgroups of meningioma. *Acta Neuropathol* 2019; 138:295–308.

Neurooncology Working Group of the German Cancer Society, Simon M, Tonn J, Stummer W, Schaub C, Weller J, Kebir S, Schäfer N, Stuplich M, Vatter H, Misch M, Keil V, Nelles M, Glas M, Coch C, Pietsch T, Hattingen E, Schmid M, Fimmers R, Weller M, Wick W, Coenen M, Urbach H, Vajkoczy P, Galldiks N, Bullinger L, Goldbrunner R, Grauer O, Krex D, Kortmann R, Hau P, Sabel M, Schlegel U, Steinbach J, Mack F, Tzaridis T, Schnell O, Bähr O, Renovanz M, Weyerbrock A, Brehmer S, Suchorska B, Schmidt-Graf F, Ringel F, Kowalski T, Tabatabai G, Seidel C, Uhl M, Herrlinger U. Lomustine–temozolamide combination therapy versus standard temozolamide therapy in patients with newly diagnosed glioblastoma with methylated MGMT promoter (CeTeG/NOA-09): a randomised, open-label, phase 3 trial. *Lancet* 2019; 393:678–688.

Katz L, Baumgarten P, Herold-Mende C, Wick W, Harter P, Weller M, von Deimling A, Snuderl M, Sen C, Wirsching H, Neidert M, Hielscher T, Liechty B, Silverman J, Zagzag D, Sen R, Wu P, Golfinos J, Reuss D, Sahm F. Loss of histone H3K27me3 identifies a subset of meningiomas with increased risk of recurrence. *Acta Neuropathol* 2018; 135:955–963.

Beier D, Proescholdt M, Reinert C, Pietsch T, Jones D, Pfister S, Hattingen E, Seidel C, Dirven L, Luerding R, Reijneveld J, Warmuth-Metz M, Bonsanto M, Bremer M, Combs S, Rieken S, Herrlinger U, Kuntze H, Mayer-Steinacker R, Moskopp D, Schneider T, Beringer A, Schlegel U, Stummer W, Welker H, Weyerbrock A, Paulsen F, Rutkowski S, Weller M, Wick W, Kortmann R, Bogdahn U, Hau P. Multicenter pilot study of radiochemotherapy as first-line treatment for adults with medulloblastoma (NOA-07). *Neuro Oncol* 2018; 20:400–410.

Wick W, Golfinopoulos V, Frenel J, Campone M, Ricard D, Marosi C, Villa S, Weyerbrock A, Hopkins K, Homicsko K, Lhermitte B, Pesce G, Roth P, Stupp R, Gorlia T, Bady P, Platten M, van den Bent M, Taphoorn M, Steuve J, Brandes A, Hamou M, Wick A, Kosch M, Weller M, Hegi M. Phase II Study of Radiotherapy and Temsirolimus versus Radiochemotherapy with Temozolomide in Patients with Newly Diagnosed Glioblastoma without MGMT Promoter Hypermethylation (EORTC 26082). *Clin Cancer Res* 2016; 22:4797–4806.

Seystahl K, Wiestler B, Hundsberger T, Happold C, Wick W, Weller M, Wick A. Bevacizumab alone or in combination with irinotecan in recurrent WHO grade II and grade III gliomas. *Eur Neurol* 2012; 69:95-101.

Hofer S, Wick W, Zander T, Weder P, Roelcke U, Pichler J, Ochsenbein A, Marosi C, Lemke D, Huber U, Hottinger A, Greil R, Elandt K, Weller M. Clinical outcome with bevacizumab in patients with recurrent high-grade glioma treated outside clinical trials. *Acta Oncol* 2011; 50:630-5.

Herrlinger U, Reifenberger G, Bamberg M, Sommer C, Tan T, Meyermann R, Wick W, Hundsberger T, Steinbach J, Kortmann R, Blaschke B, Loeser S, Koch D, Rieger J, Weller M. Phase II trial of lomustine plus temozolamide chemotherapy in addition to radiotherapy in newly diagnosed glioblastoma: UKT-03. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2006; 24:4412-7.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch