



F Harder

Contact

F Harder

Publications (6)

Langer I, Oertli D, Harder F, Wight E, Moch H, Viehl C, Ladewig A, Hsu-Schmitz S, Gueller U, Zuber M. Sentinel lymph node biopsy is associated with improved survival compared to level I & II axillary lymph node dissection in node negative breast cancer patients. *Eur J Surg Oncol* 2008; 35:805-13.

Langer I, Kocher T, Gueller U, Torhorst J, Oertli D, Harder F, Zuber M. Long-term outcomes of breast cancer patients after endoscopic axillary lymph node dissection: a prospective analysis of 52 patients. *Breast Cancer Res Treat* 2005; 90:85-91.

Bösch B, Gueller U, Schnider A, Maurer R, Harder F, Metzger U, Marti W. Perioperative detection of disseminated tumour cells is an independent prognostic factor in patients with colorectal cancer. *Br J Surg* 2003; 90:882-8.

Gueller U, Oertli D, Terracciano L, Harder F. [Neurogenic appendicopathy: a frequent, almost unknown disease picture. Evaluation of 816 appendices and review of the literature]. *Chirurg* 2001; 72:684-9.

Vogelbach P, Schütz A, Binet F, Gürke L, Gasser T, Bock A, Thiel G, Harder F. [Kidney transplantation from living donors]. *Langenbecks Archiv für Chirurgie. Supplement. Kongressband. Deutsche Gesellschaft für Chirurgie. Kongress 1997*; 114:1032-4.

Borner M, Cavalli F, Thürlimann B, Jungi W, Castiglione M, Harder F, Greiner R, Goldhirsch A, Bacchi M, Obrecht J. First isolated locoregional recurrence following mastectomy for breast cancer: results of a phase III multicenter study comparing systemic treatment with observation after excision and radiation. *Swiss Group for Clinical Cancer Research. J Clin Oncol* 1994; 12:2071-7.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch