



lan Smith

Contact

lan Smith

Publications (10)

Huober J, Hackman J, Dasappa L, Ciruelos E, Toral Pena J, Adamchuk H, Hickish T, de la Pena L, Jackisch C, Gelber R, Piccart-Gebhart M, Jouannaud C, Wildiers H, Lecocq C, Holmes E, Baselga J, de Azambuja E, Untch M, Fumagalli D, Sarp S, Láng I, Smith I, Boyle F, Xu B, Di Cosimo S. Survival outcomes of the NeoALTTO study (BIG 1-06): updated results of a randomised multicenter phase III neoadjuvant clinical trial in patients with HER2-positive primary breast cancer. Eur J Cancer 2019; 118:169-177.

Borgquist S, Regan M, Gelber R, Price K, Rabaglio M, Goldhirsch A, Coates A, Smith I, von Moos R, Ejlertsen B, Debled M, Láng I, Colleoni M, Garber J, Ahern T, Giobbie-Hurder A, Thürlimann B. Cholesterol, Cholesterol-Lowering Medication Use, and Breast Cancer Outcome in the BIG 1-98 Study. J Clin Oncol 2017; 35:1179-1188.

Chirgwin J, Colleoni M, Láng I, Neven P, Forbes J, Rabaglio M, Smith I, Goldhirsch A, Gelber R, Debled M, Ejlertsen B, Price K, Coates A, Giobbie-Hurder A, Thürlimann B. Treatment Adherence and Its Impact on Disease-Free Survival in the Breast International Group 1-98 Trial of Tamoxifen and Letrozole, Alone and in Sequence. J Clin Oncol 2016; 34:2452-9.

Regan M, BIG 1-98 Collaborative Group, Thürlimann B, Coates A, Gelber R, Price K, Rabaglio M, Wardley A, Láng I, Smith I, Forbes J, Mauriac L, Ejlertsen B, Goldhirsch A, Giobbie-Hurder A, Neven P, International Breast Cancer Study Group (IBCSG). Assessment of letrozole and tamoxifen alone and in sequence for postmenopausal women with steroid hormone receptor-positive breast cancer: the BIG 1-98 randomised clinical trial at 8·1 years median follow-up. Lancet Oncol 2011; 12:1101-8.

Chirgwin J, Coates A, Goldhirsch A, Regan M, Bonnefoi H, Ejlertsen B, Thürlimann B, Price K, Smith I, Sun Z, BIG 1-98 Collaborative and International Breast Cancer Study Groups. The advantage of letrozole over tamoxifen in the BIG 1-98 trial is consistent in younger postmenopausal women and in those with chemotherapy-induced menopause. Breast Cancer Res Treat 2011; 131:295-306.

Colleoni M, Coates A, Gelber R, Price K, Wardley A, Pienkowski T, Chirgwin J, Smith I, Láng I, Paridaens R, Forbes J, Mauriac L, Mouridsen H, Thürlimann B, Regan M, Giobbie-Hurder A, Goldhirsch A. Analyses adjusting for selective crossover show improved overall survival with adjuvant letrozole compared with tamoxifen in the BIG 1-98 study. J Clin Oncol 2011; 29:1117-24.

BIG 1-98 Collaborative Group, Gelber R, Regan M, Price K, Forbes J, Mauriac L, Smith I, Paridaens R, Thürlimann B, Goldhirsch A, Giobbie-Hurder A, Mouridsen H, Coates A. Letrozole therapy alone or in sequence with tamoxifen in women with breast cancer. NEJM 2009; 361:766-76.

Mouridsen H, Price K, Smith I, Colleoni M, Gelber R, Paridaens R, Forbes J, Mauriac L, Thürlimann B, Sun Z, Castiglione-Gertsch M, Rabaglio M, Coates A, Keshaviah A, Goldhirsch A. Cardiovascular adverse events during adjuvant endocrine therapy for early breast cancer using letrozole or tamoxifen: safety analysis of BIG 1-98 trial. JCO 2007; 25:5715-22.

Coates A, Del Mastro L, Smith I, Chirgwin J, Nogaret J, Pienkowski T, Wardley A, Jakobsen E, Price K, Láng I, Colleoni M, Keshaviah A, Thürlimann B, Mouridsen H, Mauriac L, Forbes J, Paridaens R, Castiglione-Gertsch M, Gelber R, Goldhirsch A. Five years of letrozole compared with tamoxifen as initial adjuvant therapy for postmenopausal women with endocrine-responsive early breast cancer: update of study BIG 1-98. JCO 2007; 25:486-92.

Breast International Group (BIG) 1-98 Collaborative Group, Price K, Wardly A, Wardley A, Smith I, Rabaglio M, Gelber R, Castiglione-Gertsch M, Paridaens R, Forbes J, Mauriac L, Mouridsen H, Coates A, Keshaviah A, Thürlimann B, Goldhirsch A. A comparison of letrozole and tamoxifen in postmenopausal women with early breast cancer. NEJM 2005; 353:2747-57.



No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch