



H Hawle

Contact

H Hawle

Publications (4)

Ruhstaller T, Mingrone W, Montemurro M, Girschikofsky M, Schmidt S, Bitzer M, Bedenne L, Brauchli P, Stahl M, Hess V, Mariette C, Hawle H, Thuss-Patience P, Hayoz S, Schacher S, Knorrnschild J, Schnider A, Plasswilm L, Budach W, Eisterer W, Swiss Group for Clinical Cancer Research (SAKK), the German Esophageal Cancer Study Group, the Austrian 'Arbeitsgemeinschaft Medikamentöse Tumortherapie' (AGMT), and the Fédération Francophone de Cancérologie Digestive (FFCD)/Fédération de Recherche en Ch. Neoadjuvant chemotherapy followed by chemoradiation and surgery with and without cetuximab in patients with resectable esophageal cancer: a randomized, open-label, phase III trial (SAKK 75/08). *Ann Oncol* 2018; 29:1386-1393.

Hawle H, Hess D, Mueller A, Thürlimann B. Low-Dose Fulvestrant Maintained Long-Term Complete Remission after Poor Response to Previous Endocrine Therapies in a Patient with Advanced Breast Cancer. *Case Rep Oncol* 2010; 3:131-136.

Rabaglio M, Coates A, Goldhirsch A, Gelber R, Smith I, Láng I, Nogaret J, Pienkowski T, Colleoni M, Paridaens R, Forbes J, Campone M, Mouridsen H, Thürlimann B, Hawle H, Castiglione-Gertsch M, Price K, Sun Z, BIG 1-98 Collaborative and International Breast Cancer Study Groups. Bone fractures among postmenopausal patients with endocrine-responsive early breast cancer treated with 5 years of letrozole or tamoxifen in the BIG 1-98 trial. *Ann Oncol* 2009; 20:1489-98.

Perey L, Goldhirsch A, Köberle D, Clément P, Dietrich D, Fiche M, Wildiers H, Nolé F, Zaman K, Hawle H, Paridaens R, Thürlimann B. Clinical benefit of fulvestrant in postmenopausal women with advanced breast cancer and primary or acquired resistance to aromatase inhibitors: final results of phase II Swiss Group for Clinical Cancer Research Trial (SAKK 21/00). *Ann Oncol* 2007; 18:64-9.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch