



## Heinz-Wolfram Bernd

### Contact

Heinz-Wolfram Bernd

## Publications (8)

Hartmann S, Hansmann M, Engert A, Rosenwald A, Möller P, Klapper W, Stein H, Quintanilla-Martinez L, Fend F, Cogliatti S, Ott G, Feller A, Bernd H, Mottok A, Plütschow A, Eichenauer D. The time to relapse correlates with the histopathological growth pattern in nodular lymphocyte predominant Hodgkin lymphoma. *Am J Hematol* 2019; 94:1208-1213.

Staiger A, Hartmann S, Möller P, Cogliatti S, Lenz G, Trümper L, Löffler M, Schmitz N, Pfreundschuh M, Rosenwald A, Ott G, Hansmann M, Lenze D, Ziepert M, Horn H, Scott D, Barth T, Bernd H, Feller A, Klapper W, Szczepanowski M, Hummel M, Stein H, German High-Grade Lymphoma Study Group. Clinical Impact of the Cell-of-Origin Classification and the MYC/ BCL2 Dual Expresser Status in Diffuse Large B-Cell Lymphoma Treated Within Prospective Clinical Trials of the German High-Grade Non-Hodgkin's Lymphoma Study Group. *J Clin Oncol* 2017; 35:2515-2526.

Hartmann S, Engert A, Hansmann M, Stein H, Rosenwald A, Möller P, Ott G, Feller A, Hummel M, Cogliatti S, Bernd H, Koch K, Bob R, Mottok A, Plütschow A, Eichenauer D, Klapper W. Histopathological features and their prognostic impact in nodular lymphocyte-predominant Hodgkin lymphoma--a matched pair analysis from the German Hodgkin Study Group (GHSG). *Br J Haematol* 2014; 167:238-42.

Hartmann S, Engert A, Hansmann M, Stein H, Rosenwald A, Möller P, Ott G, Feller A, Hummel M, Cogliatti S, Bernd H, Koch K, Bob R, Mottok A, Plütschow A, Eichenauer D, Klapper W. The prognostic impact of variant histology in nodular lymphocyte-predominant Hodgkin lymphoma: a report from the German Hodgkin Study Group (GHSG). *Blood* 2013; 122:4246-52; quiz 4292.

Horn H, Cogliatti S, Pfreundschuh M, Schmitz N, Trümper L, Siebert R, Loeffler M, Rosenwald A, Ott G, Möller P, Schmelter C, Hansmann M, Ziepert M, Becher C, Barth T, Bernd H, Feller A, Klapper W, Hummel M, Stein H, German High-Grade Non-Hodgkin Lymphoma Study Group. MYC status in concert with BCL2 and BCL6 expression predicts outcome in diffuse large B-cell lymphoma. *Blood* 2013; 121:2253-63.

Ott G, Frank M, Hansmann M, Barth T, Möller P, Cogliatti S, Pfreundschuh M, Schmitz N, Trümper L, Loeffler M, Müller-Hermelink H, Stein H, Ziepert M, Klapper W, Horn H, Szczepanowski M, Bernd H, Thorns C, Feller A, Lenze D, Hummel M, Rosenwald A. Immunoblastic morphology but not the immunohistochemical GCB/nonGCB classifier predicts outcome in diffuse large B-cell lymphoma in the RICOVER-60 trial of the DSHNHL. *Blood* 2010; 116:4916-25.

Bernd H, Möller P, Cogliatti S, Pfreundschuh M, Schmitz N, Trümper L, Höller S, Löffler M, Feller A, Barth T, Müller-Hermelink H, Rosenwald A, Ziepert M, Thorns C, Klapper W, Wacker H, Hummel M, Stein H, Hansmann M, Ott G, German High Grade Non-Hodgkin's Lymphoma Study Group (DSHNHL). Loss of HLA-DR expression and immunoblastic morphology predict adverse outcome in diffuse large B-cell lymphoma - analyses of cases from two prospective randomized clinical trials. *Haematologica* 2009; 94:1569-80.

Molecular Mechanisms in Malignant Lymphomas Network Project of the Deutsche Krebshilfe, Seifert M, Schübeler D, Schwaenen C, Rosenwald A, Prosper F, Pott C, Ott G, Möller P, Macleod R, Stürzenhofecker B, Weber M, Siebert R, Hasenclever D, Barker D, Esteller M, Spang R, Stein H, Trümper L, Loeffler M, Wessendorf S, Küppers R, Korn B, Ballestar E, Lopez-Serra L, Richter J, Rosolowski M, Wickham-Garcia E, Ammerpohl O, Bentink S, Bibikova M, Kreuz M, Berger H, Agirre X, Klapper W, Hummel M, Hansmann M, Fraga M, Fan J, Drexler H, Cogliatti S, Calvanese V, Bernd H, Martín-Subero J. New insights into the biology and origin of mature aggressive B-cell lymphomas by combined epigenomic, genomic, and transcriptional profiling. *Blood* 2009; 113:2488-97.

## Projects (0)

No results found.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)