



Keyur Mehta

Contact

Keyur Mehta

Publications (18)

Furlanetto J, Mehta K, Nekljudova V, Untch M, Thomssen C, Hanusch C, Lück H, Nitz U, Bauerfeind I, Gerber B, Fasching P, Huober J, Schneeweiss A, Möbus V, Lederer B, von Minckwitz G, Loibl S. Fatal events during clinical trials: an evaluation of deaths during breast cancer studies. *Breast Cancer* 2019; 26:826-834.

Denkert C, Müller V, Klauschen F, Ataseven B, Engels K, Kammler R, Pfitzner B, Diemel M, Fasching P, Colleoni M, Blohmer J, Mehta K, Budczies J, Regan M, Loibl S, Dell'Orto P, von Minckwitz G, Mastropasqua M, Solbach C, Thürlimann B, Viale G. Clinical and analytical validation of Ki-67 in 9069 patients from IBCSG VIII + IX, BIG1-98 and GeparTrio trial: systematic modulation of interobserver variance in a comprehensive in silico ring trial. *Breast Cancer Res Treat* 2019; 176:557-568.

Loibl S, Untch M, von Minckwitz G, Mehta K, Jackisch C, Tiemann K, Fasching P, Pfitzner B, Eidtmann H, Hartmann A, Lederer B, Furlanetto J, Klimowicz A, Huober J, Darb-Esfahani S, Denkert C. Integrated Analysis of PTEN and p4EBP1 Protein Expression as Predictors for pCR in HER2-Positive Breast Cancer. *Clin Cancer Res* 2016; 22:2675-83.

Loibl S, Mehta K, Nekljudova V, Blohmer J, Costa S, Denkert C, Eidtmann H, Gerber B, Hanusch C, Hilfrich J, Huober J, Schneeweiss A, Kümmel S, Paepke S, Untch M, Lederer B, Jackisch C, von Minckwitz G. Outcome after neoadjuvant chemotherapy in young breast cancer patients: a pooled analysis of individual patient data from eight prospectively randomized controlled trials. *Breast Cancer Res Treat* 2015; 152:377-87.

Fontanella C, Schneeweiss A, Paepke S, Jackisch C, Mehta K, Nekljudova V, Untch M, Neven P, von Minckwitz G, Huober J, Hilfrich J, Hanusch C, Lederer B, Gade S, Vanoppen M, Blohmer J, Costa S, Denkert C, Eidtmann H, Gerber B, Loibl S. Impact of body mass index on neoadjuvant treatment outcome: a pooled analysis of eight prospective neoadjuvant breast cancer trials. *Breast Cancer Res Treat* 2015; 150:127-39.

Klauschen F, Denkert C, Diemel M, Lederer B, Mehta K, Erbstöber E, Rüdiger T, Huober J, Blohmer J, Gerber B, Loibl S, Schmitt W, Wienert S, von Minckwitz G. Standardized Ki67 Diagnostics Using Automated Scoring--Clinical Validation in the GeparTrio Breast Cancer Study. *Clin Cancer Res* 2014; 21:3651-7.

von Minckwitz G, Huober J, Costa S, Tesch H, Hanusch C, Hilfrich J, Khandan F, Fasching P, Sinn B, Engels K, Mehta K, Nekljudova V, Klare P, Eidtmann H, Schneeweiss A, Loibl S, Salat C, Denkert C, Rezai M, Blohmer J, Jackisch C, Paepke S, Gerber B, Zahm D, Kümmel S, Untch M. Neoadjuvant carboplatin in patients with triple-negative and HER2-positive early breast cancer (GeparSixto; GBG 66): a randomised phase 2 trial. *Lancet Oncol* 2014; 15:747-56.

von Minckwitz G, Mehta K, Zahm D, Untch M, Schneeweiss A, Paepke S, Kümmel S, Kaufmann M, Jackisch C, Huober J, Hilfrich J, Hanusch C, Gerber B, Eiermann W, Eidtmann H, Denkert C, Costa S, Blohmer J, Loibl S. Response-guided neoadjuvant chemotherapy for breast cancer. *J Clin Oncol* 2013; 31:3623-30.

Denkert C, von Minckwitz G, Tesch H, Hansmann M, Reinisch M, Engels K, Sinn B, Fasching P, Mehta K, Solbach C, Brase J, Darb-Esfahani S, Kronenwett R, Prinzler J, Loibl S, Huober J, Untch M. HER2 and ESR1 mRNA expression levels and response to neoadjuvant trastuzumab plus chemotherapy in patients with primary breast cancer. *Breast Cancer Res* 2013; 15:R11.

von Minckwitz G, Mehta K, Nekljudova V, Denkert C, Konecny G, Kaufmann M, Jackisch C, Huober J, Hilfrich J, Eiermann W, Gerber B, Fasching P, Eidtmann H, Costa S, Blohmer J, Untch M, Loibl S. Definition and impact of pathologic complete response on prognosis after neoadjuvant chemotherapy in various intrinsic breast cancer subtypes. *J Clin Oncol* 2012; 30:1796-804.

von Minckwitz G, Zahm D, Khandan F, Hoffmann G, Gerber B, Eidtmann H, Fend F, Dietel M, Mehta K, du Bois A, Blohmer J, Müller B, Loibl S, Budczies J, Hanusch C, Darb-Esfahani S, Hilfrich J, Weiss E, Huober J, Denkert C. Cytoplasmic poly(adenosine diphosphate-ribose) polymerase expression is predictive and prognostic in patients with breast cancer treated with neoadjuvant chemotherapy. *J Clin Oncol* 2011; 29:2150-7.

von Minckwitz G, Gerber B, du Bois A, Huober J, Hanusch C, Konecny G, Fett W, Stickeler E, Harbeck N, Müller V, Jackisch C, Hilfrich J, Mehta K, Untch M, Nüesch E, Loibl S, Kaufmann M, Kümmel S, Fasching P, Eiermann W, Blohmer J, Costa S, Jüni P. Impact of treatment characteristics on response of different breast cancer phenotypes: pooled analysis of the German neo-adjuvant chemotherapy trials. *Breast Cancer Res Treat* 2010; 125:145-56.

Huober J, Mehta K, Darb-Esfahani S, Högel B, Thomssen C, Hauschild M, Khandan F, Belau A, Zahm D, Weiss E, Tesch H, Denkert C, von Minckwitz G, Loibl S. Effect of neoadjuvant anthracycline-taxane-based chemotherapy in different biological breast cancer phenotypes: overall results from the GeparTrio study. *Breast Cancer Res Treat* 2010; 124:133-40.

von Minckwitz G, Kühn T, du Bois A, Blohmer J, Thomssen C, Dan Costa S, Jackisch C, Kaufmann M, Mehta K, Hanusch C, Gerber B, Rezai M, Loibl S, Fasching P, Huober J, Tesch H, Bauerfeind I, Hilfrich J, Eidtmann H, Untch M. Capecitabine in addition to anthracycline- and taxane-based neoadjuvant treatment in patients with primary breast cancer: phase III GeparQuattro study. *J Clin Oncol* 2010; 28:2015-23.

Untch M, Kühn T, du Bois A, Blohmer J, Thomssen C, Dan Costa S, Jackisch C, Kaufmann M, Mehta K, Hanusch C, Gerber B, Rezai M, Loibl S, Fasching P, Huober J, Tesch H, Bauerfeind I, Hilfrich J, Eidtmann H, von Minckwitz G. Neoadjuvant treatment with trastuzumab in HER2-positive breast cancer: results from the GeparQuattro study. *J Clin Oncol* 2010; 28:2024-31.

Costa S, Tesch H, Gerber B, Baumann K, Thomssen C, Breitbach G, Ibishi S, Jackisch C, Mehta K, Weiss E, Ataseven B, Loibl S, Kaufmann M, Zahm D, Hilfrich J, Huober J, Eidtmann H, du Bois A, Blohmer J, von Minckwitz G. Neoadjuvant chemotherapy shows similar response in patients with inflammatory or locally advanced breast cancer when compared with operable breast cancer: a secondary analysis of the GeparTrio trial data. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2010; 28:83-91.

von Minckwitz G, Kaufmann M, Mehta K, Loibl S, Jackisch C, Costa S, Huober J, Gerber B, Hilfrich J, Eidtmann H, Hanusch C, Vogel P, Kümmel S, German Breast Group. Neoadjuvant vinorelbine-capecitabine versus docetaxel-doxorubicin-cyclophosphamide in early nonresponsive breast cancer: phase III randomized GeparTrio trial. *Journal of the National Cancer Institute* 2008; 100:542-51.

von Minckwitz G, Kaufmann M, Mehta K, Loibl S, Jackisch C, Costa S, Huober J, Gerber B, Hilfrich J, Eidtmann H, Hanusch C, Vogel P, Kümmel S, German Breast Group. Intensified neoadjuvant chemotherapy in early-responding breast cancer: phase III randomized GeparTrio study. *Journal of the National Cancer Institute* 2008; 100:552-62.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch