



Stig B Holmberg

Contact

Stig B Holmberg

Publications (8)

Paridaens R, Coates A, Crivellari D, Holmberg S, Price K, Thürlimann B, Gelber R, Cole B, Gelber S, Goldhirsch A. Adjuvant! Online estimation of chemotherapy effectiveness when added to ovarian function suppression plus tamoxifen for premenopausal women with estrogen-receptor-positive breast cancer. Breast Cancer Res Treat 2010; 123:303-10.

Pagani O, Goldhirsch A, Coates A, Fey M, Collins J, Thürlimann B, Lindtner J, Holmberg S, Castiglione-Gertsch M, Gelber R, Price K, International Breast Cancer Study Group (IBCSG). Patterns of recurrence of early breast cancer according to estrogen receptor status: a therapeutic target for a quarter of a century. Breast Cancer Res Treat 2009: 117:319-24.

Pagani O, Goldhirsch A, Coates A, Forbes J, Murray E, Fey M, Thürlimann B, Lindtner J, Collins J, Crivellari D, Holmberg S, Gelber R, Price K, Castiglione-Gertsch M, Simoncini E, Gelber S, International Breast Cancer Study Group. Is adjuvant chemotherapy of benefit for postmenopausal women who receive endocrine treatment for highly endocrine-responsive, node-positive breast cancer? International Breast Cancer Study Group Trials VII and 12-93. Breast Cancer Res Treat 2009: 116:491-500.

Thürlimann B, Coates A, Castiglione-Gertsch M, Forbes J, Collins J, Colleoni M, Crivellari D, Holmberg S, Gelber R, Price K, Goldhirsch A. Is chemotherapy necessary for premenopausal women with lower-risk node-positive, endocrine responsive breast cancer? 10-year update of International Breast Cancer Study Group Trial 11-93. Breast Cancer Res Treat 2009; 113:137-44.

Pestalozzi B, Goldhirsch A, Coates A, Castiglione-Gertsch M, Viale G, Murray E, Thürlimann B, Snyder R, Lindtner J, Holmberg S, Gelber R, Price K, Gusterson B, Mallon E, Zahrieh D, International Breast Cancer Study Group. Distinct clinical and prognostic features of infiltrating lobular carcinoma of the breast: combined results of 15 International Breast Cancer Study Group clinical trials. JCO 2008; 26:3006-14.

Karlsson P, Goldhirsch A, Gelber R, Forbes J, Crivellari D, Thürlimann B, Fey M, Holmberg S, Collins J, Lindtner J, Murray E, Gusterson B, Castiglione-Gertsch M, Coates A, Price K, Cole B, Wallgren A. The role of the number of uninvolved lymph nodes in predicting locoregional recurrence in breast cancer. JCO 2007; 25:2019-26.

Gruber G, Goldhirsch A, Coates A, Gelber R, Fey M, Simoncini E, Thürlimann B, Carbone A, Crivellari D, Collins J, Golouh R, Lindtner J, Holmberg S, Rudenstam C, Castiglione-Gertsch M, Price K, Nasi M, Bonetti M, International Breast Cancer Study Group. Prognostic value of extracapsular tumor spread for locoregional control in premenopausal patients with node-positive breast cancer treated with classical cyclophosphamide, methotrexate, and fluorouracil: long-term observations from International Breast Cancer Study Group Trial VI. JCO 2005; 23:7089-97.

Colleoni M, Thürlimann B, Crivellari D, Murray E, Mendiola C, Pagani O, Castiglione-Gertsch M, Coates A, Price K, Fey M, Collins J, Zahrieh D, Gelber R, Holmberg S, Mattsson J, Rudenstam C, Lindtner J, Erzen D, Snyder R, Goldhirsch A. Site of primary tumor has a prognostic role in operable breast cancer: the international breast cancer study group experience. JCO 2005; 23:1390-400.

Projects (0)

No results found.

Kantonsspital St.Gallen R

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch