



John Collins

Contact

John Collins

Publications (6)

Pagani O, Goldhirsch A, Coates A, Fey M, Collins J, Thürlimann B, Lindtner J, Holmberg S, Castiglione-Gertsch M, Gelber R, Price K, International Breast Cancer Study Group (IBCSG). Patterns of recurrence of early breast cancer according to estrogen receptor status: a therapeutic target for a quarter of a century. *Breast Cancer Res Treat* 2009; 117:319-24.

Pagani O, Goldhirsch A, Coates A, Forbes J, Murray E, Fey M, Thürlimann B, Lindtner J, Collins J, Crivellari D, Holmberg S, Gelber R, Price K, Castiglione-Gertsch M, Simoncini E, Gelber S, International Breast Cancer Study Group. Is adjuvant chemotherapy of benefit for postmenopausal women who receive endocrine treatment for highly endocrine-responsive, node-positive breast cancer? International Breast Cancer Study Group Trials VII and 12-93. *Breast Cancer Res Treat* 2009; 116:491-500.

Thürlimann B, Coates A, Castiglione-Gertsch M, Forbes J, Collins J, Colleoni M, Crivellari D, Holmberg S, Gelber R, Price K, Goldhirsch A. Is chemotherapy necessary for premenopausal women with lower-risk node-positive, endocrine responsive breast cancer? 10-year update of International Breast Cancer Study Group Trial 11-93. *Breast Cancer Res Treat* 2009; 113:137-44.

Bernhard J, Coates A, Goldhirsch A, Price K, Thürlimann B, Aebi S, Collins J, Murray E, Forbes J, Gelber R, Hürny C, Castiglione-Gertsch M, Zahrieh D, International Breast Cancer Study Group Trial VIII. Adjuvant chemotherapy followed by goserelin compared with either modality alone: the impact on amenorrhea, hot flashes, and quality of life in premenopausal patients--the International Breast Cancer Study Group Trial VIII. *JCO* 2007; 25:263-70.

Gruber G, Goldhirsch A, Coates A, Gelber R, Fey M, Simoncini E, Thürlimann B, Carbone A, Crivellari D, Collins J, Golouh R, Lindtner J, Holmberg S, Rudenstam C, Castiglione-Gertsch M, Price K, Nasi M, Bonetti M, International Breast Cancer Study Group. Prognostic value of extracapsular tumor spread for locoregional control in premenopausal patients with node-positive breast cancer treated with classical cyclophosphamide, methotrexate, and fluorouracil: long-term observations from International Breast Cancer Study Group Trial VI. *JCO* 2005; 23:7089-97.

Colleoni M, Thürlimann B, Crivellari D, Murray E, Mendiola C, Pagani O, Castiglione-Gertsch M, Coates A, Price K, Fey M, Collins J, Zahrieh D, Gelber R, Holmberg S, Mattsson J, Rudenstam C, Lindtner J, Erzen D, Snyder R, Goldhirsch A. Site of primary tumor has a prognostic role in operable breast cancer: the international breast cancer study group experience. *JCO* 2005; 23:1390-400.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch