



John F Forbes

Contact

John F Forbes

Publications (14)

Bartlett J, Rea D, Dowsett M, Cuzick J, Viale G, Forbes J, Brookes C, Van de Velde C, Putter H, Seynaeve C, Thürlimann B, Dell'Orto P, Mallon E, Sestak I, Regan M, Ahmed I. HER2 status predicts for upfront AI benefit: A TRANS-AIOG meta-analysis of 12,129 patients from ATAC, BIG 1-98 and TEAM with centrally determined HER2. Eur J Cancer 2017; 79:129-138.

Chirgwin J, Colleoni M, Láng I, Neven P, Forbes J, Rabaglio M, Smith I, Goldhirsch A, Gelber R, Debled M, Ejlertsen B, Price K, Coates A, Giobbie-Hurder A, Thürlimann B. Treatment Adherence and Its Impact on Disease-Free Survival in the Breast International Group 1–98 Trial of Tamoxifen and Letrozole, Alone and in Sequence. J Clin Oncol 2016; 34:2452-9.

Colleoni M, Coates A, Gelber R, Castiglione M, Gianni L, Thürlimann B, Forbes J, Karlsson P, Price K, Sun Z, Goldhirsch A. Annual Hazard Rates of Recurrence for Breast Cancer During 24 Years of Follow-Up: Results From the International Breast Cancer Study Group Trials I to V. J Clin Oncol 2016

Ewertz M, Goldhirsch A, Coates A, Smith I, Láng I, Colleoni M, Gelber R, Rabaglio M, Paridaens R, Forbes J, Bonnefoi H, Thürlimann B, Price K, Ejlertsen B, Regan M, Gray K, Mouridsen H. Obesity and risk of recurrence or death after adjuvant endocrine therapy with letrozole or tamoxifen in the breast international group 1–98 trial. J Clin Oncol 2012; 30:3967–75.

Regan M, BIG 1-98 Collaborative Group, Thürlimann B, Coates A, Gelber R, Price K, Rabaglio M, Wardley A, Láng I, Smith I, Forbes J, Mauriac L, Ejlertsen B, Goldhirsch A, Giobbie-Hurder A, Neven P, International Breast Cancer Study Group (IBCSG). Assessment of letrozole and tamoxifen alone and in sequence for postmenopausal women with steroid hormone receptor-positive breast cancer: the BIG 1-98 randomised clinical trial at 8·1 years median follow-up. Lancet Oncol 2011; 12:1101-8.

Colleoni M, Coates A, Gelber R, Price K, Wardley A, Pienkowski T, Chirgwin J, Smith I, Láng I, Paridaens R, Forbes J, Mauriac L, Mouridsen H, Thürlimann B, Regan M, Giobbie-Hurder A, Goldhirsch A. Analyses adjusting for selective crossover show improved overall survival with adjuvant letrozole compared with tamoxifen in the BIG 1-98 study. J Clin Oncol 2011; 29:1117-24.

Pagani O, Goldhirsch A, Coates A, Forbes J, Murray E, Fey M, Thürlimann B, Lindtner J, Collins J, Crivellari D, Holmberg S, Gelber R, Price K, Castiglione-Gertsch M, Simoncini E, Gelber S, International Breast Cancer Study Group. Is adjuvant chemotherapy of benefit for postmenopausal women who receive endocrine treatment for highly endocrine-responsive, node-positive breast cancer? International Breast Cancer Study Group Trials VII and 12-93. Breast Cancer Res Treat 2009; 116:491-500.

Thürlimann B, Coates A, Castiglione-Gertsch M, Forbes J, Collins J, Colleoni M, Crivellari D, Holmberg S, Gelber R, Price K, Goldhirsch A. Is chemotherapy necessary for premenopausal women with lower-risk node-positive, endocrine responsive breast cancer? 10-year update of International Breast Cancer Study Group Trial 11-93. Breast Cancer Res Treat 2009; 113:137-44.

Crivellari D, Chirgwin J, Smith I, Rabaglio M, Gladieff L, Del Mastro L, Láng I, Colleoni M, Gelber R, Castiglione-Gertsch M, Paridaens R, Forbes J, Mauriac L, Mouridsen H, Thürlimann B, Price K, Coates A, Sun Z, Goldhirsch A. Letrozole compared with tamoxifen for elderly patients with endocrine-responsive early breast cancer: the BIG 1-98 trial. JCO 2008; 26:1972-9.

Mouridsen H, Price K, Smith I, Colleoni M, Gelber R, Paridaens R, Forbes J, Mauriac L, Thürlimann B, Sun Z, Castiglione-Gertsch M, Rabaglio M, Coates A, Keshaviah A, Goldhirsch A. Cardiovascular adverse events during adjuvant endocrine therapy for early breast cancer using letrozole or tamoxifen: safety analysis of BIG 1-98 trial. JCO 2007; 25:5715-22.

Karlsson P, Goldhirsch A, Gelber R, Forbes J, Crivellari D, Thürlimann B, Fey M, Holmberg S, Collins J, Lindtner J, Murray E, Gusterson B, Castiglione-Gertsch M, Coates A, Price K, Cole B, Wallgren A. The role of the number of uninvolved lymph nodes in predicting locoregional recurrence in breast cancer. JCO 2007: 25:2019-26.

Coates A, Del Mastro L, Smith I, Chirgwin J, Nogaret J, Pienkowski T, Wardley A, Jakobsen E, Price K, Láng I, Colleoni M, Keshaviah A, Thürlimann B, Mouridsen H, Mauriac L, Forbes J, Paridaens R, Castiglione-Gertsch M, Gelber R, Goldhirsch A. Five years of letrozole compared with tamoxifen as initial adjuvant therapy for postmenopausal women with endocrine-responsive early breast cancer: update of study BIG 1-98. JCO 2007; 25:486-92.

Bernhard J, Coates A, Goldhirsch A, Price K, Thürlimann B, Aebi S, Collins J, Murray E, Forbes J, Gelber R, Hürny C, Castiglione-Gertsch M, Zahrieh D, International Breast Cancer Study Group Trial VIII. Adjuvant chemotherapy followed by goserelin compared with either modality alone: the impact on amenorrhea, hot flashes, and quality of life in premenopausal patients—the International Breast Cancer Study Group Trial VIII. JCO 2007; 25:263-70.

Breast International Group (BIG) 1-98 Collaborative Group, Price K, Wardly A, Wardley A, Smith I, Rabaglio M, Gelber R, Castiglione-Gertsch M, Paridaens R, Forbes J, Mauriac L, Mouridsen H, Coates A, Keshaviah A, Thürlimann B, Goldhirsch A. A comparison of letrozole and tamoxifen in postmenopausal women with early breast cancer. NEJM 2005; 353:2747-57.

Projects (0)

No results found.

Kantonsspital St.Gallen Rorschacher Strasse 95 CH-9007 St.Gallen T: +41 71 494 11 11 support.forschung@kssg.ch