



Tiziano Schepis

Contact

Tiziano Schepis

Publications (19)

Gaemperli O, Alkadhi H, Lüscher T, Eberli F, Leschka S, Scheffel H, Husmann L, Koepfli P, Valenta I, Schepis T, Kaufmann P. Functionally relevant coronary artery disease: comparison of 64-section CT angiography with myocardial perfusion SPECT. *Radiology* 2008; 248:414–23.

Husmann L, Alkadhi H, Marincek B, Leschka S, Herzog B, Desbiolles L, Stolzmann P, Hoefflinghaus T, Valenta I, Scheffel H, Schepis T, Gaemperli O, Kaufmann P. Accuracy of quantitative coronary angiography with computed tomography and its dependency on plaque composition: plaque composition and accuracy of cardiac CT. *Int J Cardiovasc Imaging* 2008; 24:895–904.

Husmann L, Alkadhi H, Marincek B, Stolzmann P, Desbiolles L, Koepfli P, Valenta I, Leschka S, Gaemperli O, Scheffel H, Schepis T, Kaufmann P. Comparison of diagnostic accuracy of 64-slice computed tomography coronary angiography in patients with low, intermediate, and high cardiovascular risk. *Acad Radiol* 2008; 15:452–61.

Husmann L, Alkadhi H, Marincek B, Stolzmann P, Desbiolles L, Leschka S, Siegrist P, Aepli U, Gaemperli O, Schepis T, Valenta I, Scheffel H, Kaufmann P. Impact of hypertension on the diagnostic accuracy of coronary angiography with computed tomography. *Int J Cardiovasc Imaging* 2008; 24:763–70.

Leschka S, Kaufmann P, Eberli F, Marincek B, Genoni M, Schepis T, Gaemperli O, Vachenaer R, Plass A, Husmann L, Koepfli P, Alkadhi H. Myocardial bridging: depiction rate and morphology at CT coronary angiography—comparison with conventional coronary angiography. *Radiology* 2008; 246:754–62.

Gaemperli O, Valenta I, Schepis T, Husmann L, Scheffel H, Desbiolles L, Leschka S, Alkadhi H, Kaufmann P. Coronary 64-slice CT angiography predicts outcome in patients with known or suspected coronary artery disease. *Eur Radiol* 2008; 18:1162–73.

Husmann L, Kaufmann P, Marincek B, Flohr T, Frauenfelder T, Cattin P, Seifert B, Gaemperli O, Schepis T, Desbiolles L, Leschka S, Alkadhi H. Coronary artery motion and cardiac phases: dependency on heart rate -- implications for CT image reconstruction. *Radiology* 2007; 245:567–76.

Schepis T, Alkadhi H, Lüscher T, Eberli F, Husmann L, Leschka S, Scheffel H, Valenta I, Namdar M, Koepfli P, Gaemperli O, Kaufmann P. Added value of coronary artery calcium score as an adjunct to gated SPECT for the evaluation of coronary artery disease in an intermediate-risk population. *J Nucl Med* 2007; 48:1424–30.

Alkadhi H, Kaufmann P, Jenni R, Marincek B, Genoni M, Flohr T, Gaemperli O, Schepis T, Vachenaer R, Scheffel H, Leschka S, Plass A, Husmann L, Desbiolles L, Frauenfelder T. Aortic regurgitation: assessment with 64-section CT. *Radiology* 2007; 245:111–21.

Gaemperli O, Alkadhi H, Husmann L, Leschka S, Desbiolles L, Stefani L, Valenta I, Namdar M, Kalff V, Schepis T, Kaufmann P. Validation of a new cardiac image fusion software for three-dimensional integration of myocardial perfusion SPECT and stand-alone 64-slice CT angiography. *Eur J Nucl Med Mol Imaging* 2007; 34:1097–106.

Gaemperli O, Alkadhi H, Husmann L, Desbiolles L, Leschka S, Soyka J, Valenta I, Koepfli P, Schepis T, Kaufmann P. Accuracy of 64-slice CT angiography for the detection of functionally relevant coronary stenoses as assessed with myocardial perfusion SPECT. *Eur J Nucl Med Mol Imaging* 2007; 34:1162–71.

Schepis T, Alkadhi H, Lüscher T, Marincek B, Wildermuth S, Eberli F, Gaemperli O, Husmann L, Desbiolles L, Leschka S, Koepfli P, Kaufmann P. Coronary artery stent geometry and in-stent contrast attenuation with 64-slice computed tomography. *Eur Radiol* 2007; 17:1464–73.

Husmann L, Kaufmann P, Marincek B, Frauenfelder T, Flohr T, Seifert B, Gaemperli O, Koepfli P, Schepis T, Desbiolles L, Leschka S, Alkadhi H. Thick maximum intensity projections for the assessment of left ventricular function with 64-slice computed tomography. *Invest Radiol* 2006; 41:746-52.

Scheffel H, Marincek B, Kaufmann P, Genoni M, Grünenfelder J, Husmann L, Schertler T, Frauenfelder T, Schepis T, Gaemperli O, Desbiolles L, Vachenauer R, Plass A, Alkadhi H, Leschka S. Accuracy of dual-source CT coronary angiography: First experience in a high pre-test probability population without heart rate control. *Eur Radiol* 2006; 16:2739-47.

Leschka S, Kaufmann P, Marincek B, Schepis T, Koepfli P, Plass A, Husmann L, Desbiolles L, Boehm T, Wildermuth S, Alkadhi H. Noninvasive coronary angiography with 64-section CT: effect of average heart rate and heart rate variability on image quality. *Radiology* 2006; 241:378-85.

Schepis T, Alkadhi H, Husmann L, Desbiolles L, Leschka S, Brunner A, Strobel K, Valenta I, Koepfli P, Gaemperli O, Kaufmann P. Comparison of 64-slice CT with gated SPECT for evaluation of left ventricular function. *J Nucl Med* 2006; 47:1288-94.

Schepis T, Alkadhi H, Husmann L, Desbiolles L, Leschka S, Burger C, Rüegg C, Koepfli P, Gaemperli O, Kaufmann P. Use of coronary calcium score scans from stand-alone multislice computed tomography for attenuation correction of myocardial perfusion SPECT. *Eur J Nucl Med Mol Imaging* 2006; 34:11-9.

Leschka S, Kaufmann P, Marincek B, Boehm T, Koepfli P, Schepis T, Gaemperli O, Desbiolles L, Husmann L, Alkadhi H. Optimal image reconstruction intervals for non-invasive coronary angiography with 64-slice CT. *Eur Radiol* 2006; 16:1964-72.

Husmann L, Kaufmann P, Marincek B, Desbiolles L, Koepfli P, Schepis T, Leschka S, Boehm T, Alkadhi H, Wildermuth S. Influence of cardiac hemodynamic parameters on coronary artery opacification with 64-slice computed tomography. *Eur Radiol* 2006; 16:1111-6.

Projects (0)

No results found.

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

support.forschung@kssg.ch