



**Ines Valenta**

**Contact**

Ines Valenta

## Publications (10)

Gaemperli O, Alkadhi H, Lüscher T, Eberli F, Leschka S, Scheffel H, Husmann L, Koepfli P, Valenta I, Schepis T, Kaufmann P. Functionally relevant coronary artery disease: comparison of 64-section CT angiography with myocardial perfusion SPECT. *Radiology* 2008; 248:414–23.

Husmann L, Alkadhi H, Marincek B, Leschka S, Herzog B, Desbiolles L, Stolzmann P, Hoefflinghaus T, Valenta I, Scheffel H, Schepis T, Gaemperli O, Kaufmann P. Accuracy of quantitative coronary angiography with computed tomography and its dependency on plaque composition: plaque composition and accuracy of cardiac CT. *Int J Cardiovasc Imaging* 2008; 24:895–904.

Husmann L, Alkadhi H, Marincek B, Stolzmann P, Desbiolles L, Koepfli P, Valenta I, Leschka S, Gaemperli O, Scheffel H, Schepis T, Kaufmann P. Comparison of diagnostic accuracy of 64-slice computed tomography coronary angiography in patients with low, intermediate, and high cardiovascular risk. *Acad Radiol* 2008; 15:452–61.

Husmann L, Alkadhi H, Marincek B, Stolzmann P, Desbiolles L, Leschka S, Siegrist P, Aepli U, Gaemperli O, Schepis T, Valenta I, Scheffel H, Kaufmann P. Impact of hypertension on the diagnostic accuracy of coronary angiography with computed tomography. *Int J Cardiovasc Imaging* 2008; 24:763–70.

Gaemperli O, Valenta I, Schepis T, Husmann L, Scheffel H, Desbiolles L, Leschka S, Alkadhi H, Kaufmann P. Coronary 64-slice CT angiography predicts outcome in patients with known or suspected coronary artery disease. *Eur Radiol* 2008; 18:1162–73.

Schepis T, Alkadhi H, Lüscher T, Eberli F, Husmann L, Leschka S, Scheffel H, Valenta I, Namdar M, Koepfli P, Gaemperli O, Kaufmann P. Added value of coronary artery calcium score as an adjunct to gated SPECT for the evaluation of coronary artery disease in an intermediate-risk population. *J Nucl Med* 2007; 48:1424–30.

Leschka S, Kaufmann P, Marincek B, Genoni M, Flohr T, Husmann L, Valenta I, Gaemperli O, Plass A, Desbiolles L, Scheffel H, Alkadhi H. Image quality and reconstruction intervals of dual-source CT coronary angiography: recommendations for ECG-pulsing windowing. *Invest Radiol* 2007; 42:543–9.

Gaemperli O, Alkadhi H, Husmann L, Leschka S, Desbiolles L, Stefani L, Valenta I, Namdar M, Kalff V, Schepis T, Kaufmann P. Validation of a new cardiac image fusion software for three-dimensional integration of myocardial perfusion SPECT and stand-alone 64-slice CT angiography. *Eur J Nucl Med Mol Imaging* 2007; 34:1097–106.

Gaemperli O, Alkadhi H, Husmann L, Desbiolles L, Leschka S, Soyka J, Valenta I, Koepfli P, Schepis T, Kaufmann P. Accuracy of 64-slice CT angiography for the detection of functionally relevant coronary stenoses as assessed with myocardial perfusion SPECT. *Eur J Nucl Med Mol Imaging* 2007; 34:1162–71.

Schepis T, Alkadhi H, Husmann L, Desbiolles L, Leschka S, Brunner A, Strobel K, Valenta I, Koepfli P, Gaemperli O, Kaufmann P. Comparison of 64-slice CT with gated SPECT for evaluation of left ventricular function. *J Nucl Med* 2006; 47:1288–94.

## Projects (0)

No results found.

---

Kantonsspital St.Gallen

Rorschacher Strasse 95

CH-9007 St.Gallen

T: +41 71 494 11 11

[support.forschung@kssg.ch](mailto:support.forschung@kssg.ch)